

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Geometrical Versions of improved Berezin-Li-Yau
Inequalities

Leander Geisinger, Ari Laptev, Timo Weidl

Preprint 2010/009

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Geometrical Versions of improved Berezin-Li-Yau
Inequalities

Leander Geisinger, Ari Laptev, Timo Weidl

Preprint 2010/009

Fachbereich Mathematik
Fakultät Mathematik und Physik
Universität Stuttgart
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de

WWW: <http://www.mathematik.uni-stuttgart.de/preprints>

ISSN **1613-8309**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.

LaTeX-Style: Winfried Geis, Thomas Merkle

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^d$ be an open set and let $-\Delta$ denote the Dirichlet Laplace operator on $L^2(\Omega)$, defined as a self-adjoint operator with form domain $H_0^1(\Omega)$. We assume that the volume of Ω , denoted by $|\Omega|$, is finite. Then the embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact and the spectrum of $-\Delta$ is discrete: It consists of positive eigenvalues

$$0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$$

accumulating at infinity only.

Here we are interested in upper bounds on the Riesz means

$$\sum_k (\Lambda - \lambda_k)_+^\sigma = \text{Tr}(-\Delta - \Lambda)_-^\sigma, \quad \sigma \geq 0,$$

where we use the notation $x_\pm = (|x| \pm x)/2$. In 1972 Berezin proved that, convex eigenvalue means are bounded uniformly by the corresponding phase-space volume, see [Ber72]: For any open set $\Omega \subset \mathbb{R}^d$, $\sigma \geq 1$, and all $\Lambda > 0$

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma \leq \frac{1}{(2\pi)^d} \iint_{\Omega \times \mathbb{R}^d} (|p|^2 - \Lambda)_+^\sigma dp dx = L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2}, \quad (1.1)$$

see also [LY83], where the problem is treated from a different point of view. Here $L_{\sigma,d}^{cl}$ denotes the so-called Lieb-Thirring constant

$$L_{\sigma,d}^{cl} = \frac{\Gamma(\sigma + 1)}{(4\pi)^{d/2} \Gamma(\sigma + 1 + d/2)}.$$

The Berezin inequality (1.1) captures, in particular, the well-known asymptotic limit that goes back to Hermann Weyl [Wey12]: For $\Omega \subset \mathbb{R}^d$ and $\sigma \geq 0$ the asymptotic identity

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma = L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2} + o\left(\Lambda^{\sigma+d/2}\right) \quad (1.2)$$

holds true as $\Lambda \rightarrow \infty$. From this follows, that the Berezin inequality is sharp, in the sense that the constant in (1.1) cannot be improved. However, Hermann Weyl's work stimulated further analysis of the asymptotic formula and (1.2) was gradually improved by studying the second term, see [CH24, Hör68, Iv80, Mel80, SV97, Iv98] and references within. The precise second term was found by Ivrii [Ivr80]: Under appropriate conditions on the set Ω and its boundary $\partial\Omega$ the relation

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma = L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2} - \frac{1}{4} L_{\sigma,d-1}^{cl} |\partial\Omega| \Lambda^{\sigma+(d-1)/2} + o\left(\Lambda^{\sigma+(d-1)/2}\right) \quad (1.3)$$

holds as $\Lambda \rightarrow \infty$. To simplify notation we write $|\Omega|$ for the volume (the d -dimensional Lebesgue measure) of Ω , as well as $|\partial\Omega|$ for the $d-1$ -dimensional surface area of its boundary. Since the second term of this semi-classical limit is negative, the question arises, whether the Berezin inequality (1.1) can be improved by a negative remainder term.

Recently, several such improvements have been found, initially for the discrete Laplace operator, see [FLU02]. The first result for the continuous Laplace operator is due to Melas [Mel03]. From his work follows that

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma,d}^{cl} |\Omega| \left(\Lambda - M_d \frac{|\Omega|}{I(\Omega)} \right)_+^{\sigma+d/2}, \quad \Lambda > 0, \quad \sigma \geq 1, \quad (1.4)$$

where M_d is a constant depending only on the dimension and $I(\Omega)$ denotes the second moment of Ω , see also [Ily09, Yol09] for further generalisations. One should mention, however, that

these corrections do not capture the correct order in Λ from the second term of the asymptotics (1.3). This was improved in the two-dimensional case in [KVV09], where it is shown that one can choose the order of the correction term arbitrarily close to the correct one.

In this paper we are interested in the case $\sigma \geq 3/2$. For these values of σ it is known, [Wei08], that one can strengthen the Berezin inequality for any open set $\Omega \subset \mathbb{R}^d$ with a negative remainder term reflecting the correct order in Λ in comparison to the second term of (1.3). However, since one can increase $|\partial\Omega|$ without changing the individual eigenvalues λ_k significantly, a direct analog of the first two terms of the asymptotics (1.3) cannot yield a uniform bound on the eigenvalue means. Therefore - without further conditions on Ω - any uniform improvement of (1.1) must invoke other geometric quantities.

In the result from [Wei08] the remainder term involves certain projections on $d - 1$ -dimensional hyperplanes. In [GW10] a universal improvement of (1.1) was found, that holds for $\sigma \geq 3/2$ with a correction term of correct order, depending only on the volume of Ω .

The proof of the aforementioned results relies on operator-valued Lieb-Thirring inequalities [LW00] and an inductive argument, that allows to reduce the problem to estimating traces of the one-dimensional Dirichlet Laplace operator on open intervals.

In this paper we use the same approach, but with new estimates in the one-dimensional case, in order to make the dependence on the geometry more transparent. The new one-dimensional bounds involve the distance to the boundary of the interval in question and are related to Hardy-Lieb-Thirring inequalities for Schrödinger operators, see [EF06]. There it is shown that for $\sigma \geq 1/2$ and potentials $V \in L^{\sigma+1/2}(\mathbb{R}_+)$, given on the half-line $\mathbb{R}_+ = (0, \infty)$, the inequality

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - V \right)_-^\sigma \leq L_\sigma \int_0^\infty \left(V(t) - \frac{1}{4t^2} \right)_+^{\sigma+1/2} dt$$

holds true, with a constant L_σ independent of V . For further developments see [FLS08, Fra09].

We start this paper with analysing the special case of the Dirichlet Laplace operator given on a finite interval $I \subset \mathbb{R}$, with the constant potential $V \equiv \Lambda$. For $\sigma \geq 1$ we establish that the estimate

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_-^\sigma \leq L_{\sigma,1}^{\mathrm{cl}} \int_I \left(\Lambda - \frac{1}{4\delta(t)^2} \right)_+^{\sigma+1/2} dt$$

is valid with the sharp constant $L_{\sigma,1}^{\mathrm{cl}}$, where $\delta(t)$ denotes the distance to the boundary of I . This is done in section 2.

Then we can use results from [LW00, Wei08] to deduce bounds in higher dimensions: In section 3 we first derive improvements of (1.1), which are valid for any open set $\Omega \subset \mathbb{R}^d$, $d \geq 2$. These improvements depend on the geometry of Ω . In view of the asymptotic result (1.3) one might expect, that this geometric dependence can be expressed in terms of the boundary of Ω . To see this, we adapt methods, which were used in [Dav94, Dav99, HHL02] to derive geometric versions of Hardy's inequality. Here the result gives an improved Berezin inequality with a correction term of correct order depending on geometric properties of the boundary.

If Ω is convex and smooth this dependence can be expressed only in terms of $|\Omega|$, $|\partial\Omega|$ and the curvature of the boundary. In particular the first remainder term of the estimate is very similar to the second term of the semiclassical asymptotics (1.3): it shows the same order in Λ and it depends only on the surface area of the boundary.

In section 4 we return to the general case, where $\Omega \subset \mathbb{R}^d$ is not necessarily convex or smooth, and obtain lower bounds on individual eigenvalues λ_k . Under certain conditions on

the geometry of Ω these results improve the estimate

$$\lambda_k \geq C_d \frac{d}{d+2} \left(\frac{k}{|\Omega|} \right)^{2/d}, \quad (1.5)$$

from [LY83], where C_d denotes the semi-classical constant $4\pi \Gamma(d/2 + 1)^{2/d}$.

Finally, in section 5, we specialise to the two-dimensional case, where we can use the foregoing results and refined geometric considerations to further improve and generalise the inequalities. In particular, we avoid dependence on curvature, thus we do not require smoothness of the boundary.

The question whether such improved estimates can be generalised to $1 \leq \sigma < 3/2$ remains open.

2. ONE-DIMENSIONAL CONSIDERATIONS

Let us consider an open interval $I \subset \mathbb{R}$ of length $l > 0$. For $t \in I$ let

$$\delta(t) = \inf \{|t - s| : s \notin I\}$$

be the distance to the boundary. The eigenvalues of $-d^2/dt^2$ subject to Dirichlet boundary conditions at the endpoints of I are given by $\lambda_k = k^2\pi^2/l^2$. Therefore the Riesz means equal

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_-^\sigma = \sum_k \left(\Lambda - \frac{k^2\pi^2}{l^2} \right)_+^\sigma.$$

To find precise bounds on the Riesz means in the one-dimensional case, it suffices to analyse this sum explicitly. Our main observation is

Lemma 1. *Let $I \subset \mathbb{R}$ be an open interval and $\sigma \geq 1$. Then the estimate*

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_-^\sigma \leq L_{\sigma,1}^{cl} \int_I \left(\Lambda - \frac{1}{4\delta(t)^2} \right)_+^{\sigma+1/2} dt,$$

holds true for all $\Lambda > 0$. The constant $1/4$ cannot be improved.

The remainder of this section deals with the proof of this estimate. First we need two rather technical results, whose proof is elementary but not trivial and therefore will be given in the appendix.

Lemma 2. *For all $A \geq 1/\pi$*

$$\sum_k \left(1 - \frac{k^2}{A^2} \right)_+ \leq \frac{2}{3\pi} \int_1^{\pi A} \left(1 - \frac{1}{s^2} \right)^{3/2} ds. \quad (2.1)$$

Lemma 3. *Let $I \subset \mathbb{R}$ be an open interval of length $l > 0$. Then for $\sigma \geq 1$ and $c > 0$*

$$L_{\sigma,1}^{cl} \int_I \left(\Lambda - \frac{c}{\delta(t)^2} \right)_+^{\sigma+1/2} dt - \sum_k \left(\Lambda - \frac{\pi^2 k^2}{l^2} \right)_+^\sigma = \left(\frac{1}{2} - \sqrt{c} \right) \Lambda^\sigma + o(\Lambda^\sigma)$$

holds as $\Lambda \rightarrow \infty$.

Proof of Lemma 1. Note that one can always assume $I = (0, l)$, where $l > 0$ denotes the length of the interval. First we deduce the estimate for $\sigma = 1$ from Lemma 2. Assume $\Lambda \geq l^{-2}$ and apply Lemma 2 with $A = l\sqrt{\Lambda}/\pi$ to get

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_- = \Lambda \sum_k \left(1 - \frac{\pi^2 k^2}{l^2 \Lambda} \right)_+ \leq \Lambda \frac{2}{3\pi} \int_1^{l\sqrt{\Lambda}} \left(1 - \frac{1}{s^2} \right)^{3/2} ds.$$

Substituting $s = 2t\sqrt{\Lambda}$, we find that

$$\mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_- \leq \frac{4}{3\pi} \int_{1/(2\sqrt{\Lambda})}^{l/2} \left(\Lambda - \frac{1}{4t^2} \right)_+^{3/2} dt = 2L_{1,1}^{cl} \int_0^{l/2} \left(\Lambda - \frac{1}{4t^2} \right)_+^{3/2} dt$$

holds for all $\Lambda \geq l^{-2}$. Note that this inequality is trivially true for $0 < \Lambda < l^{-2}$, since the left hand side is zero. Finally, we use the identities

$$\int_0^{l/2} \left(\Lambda - \frac{1}{4t^2} \right)_+^{3/2} dt = \int_{l/2}^l \left(\Lambda - \frac{1}{4(l-t)^2} \right)_+^{3/2} dt = \frac{1}{2} \int_0^l \left(\Lambda - \frac{1}{4\delta(t)^2} \right)_+^{3/2} dt$$

to finish the proof for $\sigma = 1$.

To deduce the claim for $\sigma > 1$ we can apply a method from [AL78]. Writing

$$\sum_k (\Lambda - \lambda_k)_+^\sigma = \frac{1}{B(2, \sigma - 1)} \int_0^\infty \tau^{\sigma-2} \sum_k (\Lambda - \tau - \lambda_k)_+ d\tau,$$

we estimate

$$\begin{aligned} \mathrm{Tr} \left(-\frac{d^2}{dt^2} - \Lambda \right)_-^\sigma &\leq \frac{1}{B(2, \sigma - 1)} L_{1,1}^{cl} \int_I \int_0^\infty \tau^{\sigma-2} \left(\Lambda - \frac{1}{4\delta(t)^2} - \tau \right)_+^{3/2} d\tau dt \\ &= L_{1,1}^{cl} \frac{B(5/2, \sigma - 1)}{B(2, \sigma - 1)} \int_I \left(\Lambda - \frac{1}{4\delta(t)^2} \right)_+^{\sigma+1/2} dt \end{aligned}$$

and the result follows from the identity $L_{1,1}^{cl} B(5/2, \sigma - 1) = L_{\sigma,1}^{cl} B(2, \sigma - 1)$.

The claim, that the constant $1/4$ cannot be improved, follows from Lemma 3: For $c = 1/4$ the leading term of the asymptotics in Lemma 3 vanishes. For any constant $c > 1/4$ the leading term is negative, thus the estimate from Lemma 1 must fail in this case, for large values of Λ . \square

Figure 1 illustrates the results of Lemma 1 and Lemma 3 for $l = \pi^2$ and $\sigma = 1$ with the sharp constant $c = 1/4$: The function

$$f(\Lambda) = L_{1,1}^{cl} \int_0^\pi \left(\Lambda - \frac{1}{4\delta(t)^2} \right)_+^{3/2} dt - \sum_k (\Lambda - k^2)_+$$

is plotted for $1 < \Lambda < 112$, so that the first ten minima are shown.

3. RESULTS IN HIGHER DIMENSIONS

In this section we use the one-dimensional result to prove uniform eigenvalue estimates for the Dirichlet Laplace operator in bounded open sets in higher dimensions. These estimates - refinements of the Berezin inequality (1.1) - depend on the geometry of the set, in particular on properties of the boundary.




FIGURE 1. The function $f(\Lambda)$, illustrating the results from section 2.

3.1. Arbitrary open sets. First we provide general estimates, valid for any open subset $\Omega \subset \mathbb{R}^d$, $d \geq 2$. Let

$$\mathbb{S}^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$$

denote the unit-sphere in \mathbb{R}^d . For an arbitrary direction $u \in \mathbb{S}^{d-1}$ and $x \in \Omega$ set

$$\begin{aligned} \theta(x, u) &= \inf \{t > 0 : x + tu \notin \Omega\}, \\ d(x, u) &= \inf \{\theta(x, u), \theta(x, -u)\} \quad \text{and} \\ l(x, u) &= \theta(x, u) + \theta(x, -u). \end{aligned}$$

Theorem 4. *Let $\Omega \subset \mathbb{R}^d$ be an open set and let $u \in \mathbb{S}^{d-1}$ and $\sigma \geq 3/2$. Then for all $\Lambda > 0$ the estimate*

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma, d}^{cl} \int_{\Omega} \left(\Lambda - \frac{1}{4d(x, u)^2} \right)_+^{\sigma+d/2} dx \quad (3.1)$$

holds true.

Remark. Let us define

$$l_0 = \inf_{u \in \mathbb{S}^{d-1}} \sup_{x \in \Omega} l(x, u). \quad (3.2)$$

Then Theorem 4 implies the following improvement of Melas-type: For $\sigma \geq 3/2$ and all $\Lambda > 0$ the estimate

$$\text{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma, d}^{cl} |\Omega| \left(\Lambda - \frac{1}{l_0^2} \right)_+^{\sigma+d/2} \quad (3.3)$$

holds. In convex domains l_0 is the minimal width of the domain, see [BF48]. In this case $1/l_0^2$ is bounded from below by a multiple of $|\Omega|^{-2/d}$, [YB61], while no such bound holds for the improving term $|\Omega|/I(\Omega)$ in Melas' inequality (1.4).

The proof of Theorem 4 relies on a lifting technique, which was introduced in [Lap97], see also [LW00, ELW04, Wei08, FL08] for further developments and applications.

Proof of Theorem 4. We apply the argument used in [Wei08] to reduce the problem to one-dimensional estimates. Fix a Cartesian coordinate system in \mathbb{R}^d , such that the given direction u corresponds to the vector $(0, \dots, 0, 1)$.

For $x \in \mathbb{R}^d$ write $x = (x', t) \in \mathbb{R}^{d-1} \times \mathbb{R}$ and let ∇' and $-\Delta'$ denote the gradient and the Laplace operator in the first $d-1$ dimensions. Each section $\Omega(x') = \{t \in \mathbb{R} : (x', t) \in \Omega\}$ consists of at most countably many open intervals $J_k(x') \subset \mathbb{R}$, $k = 1, \dots, N(x') \leq \infty$.

We consider the quadratic form of $-\Delta - \Lambda$ on functions φ from the form core $C_0^\infty(\Omega)$ and write

$$\|\nabla\varphi\|_{L^2(\Omega)}^2 - \Lambda \|\varphi\|_{L^2(\Omega)}^2 = \|\nabla'\varphi\|_{L^2(\Omega)}^2 + \int_{\mathbb{R}^{d-1}} dx' \int_{\Omega(x')} \left(|\partial_t \varphi|^2 - \Lambda |\varphi|^2 \right) dt.$$

The functions $\varphi(x', \cdot)$ satisfy Dirichlet boundary conditions at the endpoints of each interval $J_k(x')$ forming $\Omega(x')$, hence

$$\begin{aligned} \int_{\Omega(x')} \left(|\partial_t \varphi|^2 - \Lambda |\varphi|^2 \right) dt &= \sum_{k=1}^{N(x')} \int_{J_k(x')} \left(|\partial_t \varphi|^2 - \Lambda |\varphi|^2 \right) dt \\ &\geq - \sum_{k=1}^{N(x')} \langle V_k(x') \varphi(x', \cdot), \varphi(x', \cdot) \rangle_{L^2(J_k(x'))}, \end{aligned}$$

where the bounded, non-negative operators $V_k(x', \Lambda) = (-\partial_t^2 - \Lambda)_-$ are the negative parts of the Sturm-Liouville operators $-\partial_t^2 - \Lambda$ with Dirichlet boundary conditions on $J_k(x')$. Let

$$V(x', \Lambda) = \bigoplus_{k=1}^{N(x')} V_k(x', \Lambda)$$

be the negative part of $-\partial_t^2 - \Lambda$ on $\Omega(x')$ subject to Dirichlet boundary conditions on the endpoints of each interval $J_k(x')$, $k = 1, \dots, N(x')$, that is on $\partial\Omega(x')$. Then

$$\int_{\Omega(x')} \left(|\partial_t \varphi|^2 - \Lambda |\varphi|^2 \right) dt \geq - \langle V(x', \Lambda) \varphi(x', \cdot), \varphi(x', \cdot) \rangle_{L^2(\Omega(x'))}$$

and consequently

$$\|\nabla\varphi\|_{L^2(\Omega)}^2 - \Lambda \|\varphi\|_{L^2(\Omega)}^2 \geq \|\nabla'\varphi\|_{L^2(\Omega)}^2 - \int_{\mathbb{R}^{d-1}} dx' \langle V \varphi(x', \cdot), \varphi(x', \cdot) \rangle_{L^2(\Omega(x'))}.$$

Now we can extend this quadratic form by zero to $C_0^\infty(\mathbb{R}^d \setminus \partial\Omega)$, which is a form core for $(-\Delta_{\mathbb{R}^d \setminus \Omega}) \oplus (-\Delta_\Omega - \Lambda)$. This operator corresponds to the left hand side of the equality above, while the semi-bounded form on the right hand side is closed on the larger domain $H^1(\mathbb{R}^{d-1}, L^2(\mathbb{R}))$, where it corresponds to the operator

$$-\Delta' \otimes \mathbb{I} - V(x', \Lambda) \quad \text{on} \quad L^2(\mathbb{R}^{d-1}, L^2(\mathbb{R})). \quad (3.4)$$

Due to the positivity of $-\Delta_{\mathbb{R}^d \setminus \Omega}$ we can use the variational principle to deduce that for any $\sigma \geq 0$

$$\begin{aligned} \text{Tr}(-\Delta_\Omega - \Lambda)_-^\sigma &= \text{Tr} \left(\left(-\Delta_{\mathbb{R}^d \setminus \Omega} \right) \oplus (-\Delta_\Omega - \Lambda) \right)_-^\sigma \\ &\leq \text{Tr}(-\Delta' \otimes \mathbb{I} - V(x', \Lambda))_-^\sigma. \end{aligned}$$

Now we apply sharp Lieb-Thirring inequalities [LW00] to the Schrödinger operator (3.4) with the operator-valued potential $-V(x', \Lambda)$ and obtain that for $\sigma \geq 3/2$

$$\mathrm{Tr}(-\Delta_\Omega - \Lambda)_-^\sigma \leq L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \mathrm{Tr} V(x', \Lambda)^{\sigma+(d-1)/2} dx'. \quad (3.5)$$

To estimate the trace of the one-dimensional differential operator $V(x', \Lambda)$ we apply Lemma 1. Our choice of coordinate system implies that for $x = (x', t) \in J_k(x')$ the distance of t to the boundary of the interval $J_k(x')$ is given by $d(x, u)$. Hence, Lemma 1 implies

$$\begin{aligned} \mathrm{Tr} V(x', \Lambda)^{\sigma+(d-1)/2} &= \sum_{k=1}^{N(x')} \mathrm{Tr} \left(-\frac{d^2}{dt^2} \Big|_{J_k(x')} - \Lambda \right)_-^{\sigma+(d-1)/2} \\ &\leq L_{\sigma+(d-1)/2, 1}^{cl} \int_{\Omega(x')} \left(\Lambda - \frac{1}{4d((x', t), u)} \right)_+^{\sigma+d/2} dt \end{aligned}$$

and the result follows from (3.5) and the identity $L_{\sigma, d-1}^{cl} L_{\sigma+(d-1)/2, 1}^{cl} = L_{\sigma, d}^{cl}$. \square

We proceed to analysing the geometric properties of (3.1). Note that the left hand side of (3.1) is independent of the choice of direction $u \in \mathbb{S}^{d-1}$, while the right hand side depends on u and therefore on the geometry of Ω . For a given set Ω one can minimise the right hand side in $u \in \mathbb{S}^{d-1}$. The result, however, depends on the geometry of Ω in a rather tricky way. In order to make this geometric dependence more transparent, we average the right hand side of (3.1) over $u \in \mathbb{S}^{d-1}$. Even though the resulting bound is - in general - not as precise as (3.1), it allows a more appropriate geometric interpretation.

To analyse the effect of the boundary, one would like to estimate $d(x, u)$ in terms of the distance to the boundary, see [Dav94, Dav99, HHL02], where this approach is used to derive geometrical versions of Hardy's inequality. To avoid complications that arise, for example, if the complement of Ω contains isolated points, we use slightly different notions: For $x \in \Omega$ let

$$\Omega(x) = \{y \in \Omega : x + t(y - x) \in \Omega, \forall t \in [0, 1]\}$$

be the part of Ω that "can be seen" from x and let

$$\delta(x) = \inf \left\{ |y - x| : y \notin \overline{\Omega(x)} \right\}$$

denote the distance to the exterior of $\Omega(x)$.

For fixed $\varepsilon > 0$ put

$$A_\varepsilon(x) = \left\{ a \in \mathbb{R}^d \setminus \overline{\Omega(x)} : |x - a| < \delta(x) + \varepsilon \right\}$$

and for $a \in A_\varepsilon(x)$ set $B_x(a) = \{y \in \mathbb{R}^d : |y - a| < |x - a|\}$ and

$$\rho_a(x) = \frac{|B_x(a) \setminus \overline{\Omega(x)}|}{\omega_d |x - a|^d},$$

where ω_d denotes the volume of the unit ball in \mathbb{R}^d . To get a result, independent of a and ε , set

$$\rho(x) = \inf_{\varepsilon > 0} \sup_{a \in A_\varepsilon(x)} \rho_a(x).$$

Note that $\mathbb{R}^d \setminus \overline{\Omega(x)}$ is open, hence $\rho_a(x) > 0$ and $\rho(x) > 0$ hold for $x \in \Omega$. Finally, we define

$$M_\Omega(\Lambda) = \int_{R_\Omega(\Lambda)} \rho(x) dx,$$

where $R_\Omega(\Lambda) \subset \Omega$ denotes the set $\{x \in \Omega : \delta(x) < 1/(4\sqrt{\Lambda})\}$. The main result of this section allows a geometric interpretation of the remainder term:

Theorem 5. *Let $\Omega \subset \mathbb{R}^d$ be an open set with finite volume and $\sigma \geq 3/2$. Then for all $\Lambda > 0$ we have*

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2} - L_{\sigma,d}^{cl} 2^{-d+1} \Lambda^{\sigma+d/2} M_\Omega(\Lambda). \quad (3.6)$$

The function $\rho(x)$ depends on the behaviour of the boundary close to $x \in \Omega$. For example, $\rho(x)$ is small close to a cusp. On the other hand $\rho(x)$ is larger than $1/2$ in a convex domain. By definition, the function $M_\Omega(\Lambda)$ gives an average of this behaviour over $R_\Omega(\Lambda)$, which is like a tube of width $1/(4\sqrt{\Lambda})$ around the boundary.

Note that $M_\Omega(\Lambda)$ tends to zero as Λ tends to infinity. This decay in Λ is of the order $(\delta_M - d)/2$, where δ_M denotes the interior Minkowski dimension of the boundary, see e.g. [Lap91, FV93, FLV95] for definition and examples. If $d - 1 \leq \delta_M < d$ and if the upper Minkowski content of the boundary is finite, then the second term of the asymptotic limit of the Riesz means equals $O(\Lambda^{\sigma+\delta_M/2})$ as $\Lambda \rightarrow \infty$, see [Lap91]. Therefore the remainder term in (3.6) reflects the correct order of growth in the asymptotic limit.

In particular, if the dimension of the boundary equals $d - 1$, we find

$$M_\Omega(\Lambda) = |\partial\Omega| \Lambda^{-1/2} + o(\Lambda^{-1/2})$$

as $\Lambda \rightarrow \infty$ and the second term in (3.6) is in close correspondence with the asymptotic formula (1.3).

Proof of Theorem 5. We start from the result of Theorem 4 and average over all directions to get

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma,d}^{cl} \Lambda^{\sigma+d/2} \int_\Omega \int_{\mathbb{S}^{d-1}} \left(1 - \frac{1}{4\Lambda d(x,u)^2}\right)_+^{\sigma+d/2} d\nu(u) dx, \quad (3.7)$$

where $d\nu(u)$ denotes the normed measure on \mathbb{S}^{d-1} .

For $x \in \Omega$ and $a \notin \overline{\Omega(x)}$ let $\Theta(x,a) \subset \mathbb{S}^{d-1}$ be the subset of all directions $u \in \mathbb{S}^{d-1}$, satisfying $x + su \in B_x(a) \setminus \overline{\Omega(x)}$ for some $s > 0$. For such s we have

$$s \leq 2|x - a|. \quad (3.8)$$

By definition of $\rho_a(x)$ and $\Theta(x,a)$ we find

$$\rho_a(x) \omega_d |x - a|^d = |B_x(a) \setminus \overline{\Omega(x)}| \leq \int_{\Theta(x,a)} d\nu(u) \omega_d (2|x - a|)^d,$$

hence

$$\int_{\Theta(x,a)} d\nu(u) \geq 2^{-d} \rho_a(x). \quad (3.9)$$

Using (3.8) we also see that for $u \in \Theta(x,a)$ the estimate $d(x,u) \leq s \leq 2|x - a|$ holds.

Now fix $\Lambda > 0$ and choose $0 < \varepsilon < 1/(4\sqrt{\Lambda})$ and $a \in A_\varepsilon(x)$. By definition of $A_\varepsilon(x)$ it follows that for all $u \in \Theta(x,a)$

$$d(x,u) \leq 2|x - a| < 2(\delta(x) + \varepsilon). \quad (3.10)$$

The set $\Theta(x, a)$ must be contained in one hemisphere of \mathbb{S}^{d-1} which we denote by \mathbb{S}_+^{d-1} . Using that $d(x, u) = d(x, -u)$ we estimate

$$\begin{aligned} \int_{\mathbb{S}^{d-1}} \left(1 - \frac{1}{4\Lambda d(x, u)^2}\right)_+^{\sigma+d/2} d\nu(u) &= 2 \int_{\mathbb{S}_+^{d-1}} \left(1 - \frac{1}{4\Lambda d(x, u)^2}\right)_+^{\sigma+d/2} d\nu(u) \\ &\leq 1 - 2 \int_{\{u \in \mathbb{S}_+^{d-1} : d(x, u) \leq 1/(2\sqrt{\Lambda})\}} d\nu(u). \end{aligned}$$

Assume that $\delta(x) \leq 1/(4\sqrt{\Lambda}) - \varepsilon$. From (3.10) it follows that

$$\Theta(x, a) \subset \left\{u \in \mathbb{S}_+^{d-1} : d(x, u) \leq 1/(2\sqrt{\Lambda})\right\},$$

hence, using (3.9), we conclude

$$\int_{\mathbb{S}^{d-1}} \left(1 - \frac{1}{4\Lambda d(x, u)^2}\right)_+^{\sigma+d/2} d\nu(u) \leq 1 - 2 \int_{\Theta(x, a)} d\nu(u) \leq 1 - 2^{1-d} \rho_a(x).$$

Since $a \in A_\varepsilon(x)$ was arbitrary we arrive at

$$\int_{\mathbb{S}^{d-1}} \left(1 - \frac{1}{4\Lambda d(x, u)^2}\right)_+^{\sigma+d/2} d\nu(u) \leq 1 - 2^{1-d} \rho(x),$$

for all $x \in \Omega$ with $\delta(x) \leq 1/(4\sqrt{\Lambda}) - \varepsilon$ and we can take the limit $\varepsilon \rightarrow 0$.

It follows that

$$\int_{\Omega} \int_{\mathbb{S}^{d-1}} \left(1 - \frac{1}{4\Lambda d(x, u)^2}\right)_+^{\sigma+d/2} d\nu(u) dx \leq |\Omega| - 2^{1-d} \int_{\{x \in \Omega : \delta(x) < 1/(4\sqrt{\Lambda})\}} \rho(x) dx$$

and inserting this into (3.7) yields the claimed result. \square

3.2. Convex domains. If $\Omega \subset \mathbb{R}^d$ is convex, we have $\Omega(x) = \Omega$ and

$$\rho(x) > 1/2 \tag{3.11}$$

for all $x \in \Omega$. Thus we can simplify the remainder term, by estimating $M_\Omega(\Lambda)$.

Corollary 6. *Let $\Omega \subset \mathbb{R}^d$ be a bounded, convex domain with smooth boundary and assume that the curvature of $\partial\Omega$ is bounded from above by $1/R$. Then for $\sigma \geq 3/2$ and all $\Lambda > 0$ we have*

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma, d}^{cl} |\Omega| \Lambda^{\sigma+d/2} - L_{\sigma, d}^{cl} 2^{-d-2} |\partial\Omega| \Lambda^{\sigma+(d-1)/2} \int_0^1 \left(1 - \frac{d-1}{4R\sqrt{\Lambda}} s\right)_+ ds.$$

Proof. Inserting (3.11) into the definition of $M_\Omega(\Lambda)$ yields

$$M_\Omega(\Lambda) > \frac{1}{2} \int_{R_\Omega(\Lambda)} dx.$$

Let $\Omega_t = \{x \in \Omega : \delta(x) > t\}$ be the inner parallel set of Ω and write

$$\int_{R_\Omega(\Lambda)} dx = \int_{\{x \in \Omega : \delta(x) < 1/(4\sqrt{\Lambda})\}} dx = \int_0^{1/(4\sqrt{\Lambda})} |\partial\Omega_t| dt.$$

Now we can use Steiner's Theorem, see [Gug77, vdB84], namely

$$|\partial\Omega_t| \geq \left(1 - \frac{d-1}{R}t\right)_+ |\partial\Omega|. \quad (3.12)$$

It follows that

$$M_\Omega(\Lambda) > \frac{1}{2}|\partial\Omega| \int_0^{1/(4\sqrt{\Lambda})} \left(1 - \frac{d-1}{R}t\right)_+ dt = \frac{|\partial\Omega|}{8\sqrt{\Lambda}} \int_0^1 \left(1 - \frac{d-1}{4R\sqrt{\Lambda}}s\right)_+ ds.$$

Inserting this into (3.6) completes the proof. \square

Let us single out the case where $\Omega = B_r$ is a ball in \mathbb{R}^d with radius $r > 0$. Note that the first eigenvalue of the Dirichlet Laplace operator on B_r is given by

$$\lambda_1(B_r) = \frac{\pi j_{d/2-1,1}^2}{\Gamma(d/2+1)^{2/d} |B_r|^{2/d}},$$

where $j_{d/2-1,1}$ denotes the first zero of the Bessel function $J_{d/2-1}$. Again, we have to estimate

$$M_\Omega(\Lambda) > \frac{1}{2} \int_0^{1/(4\sqrt{\Lambda})} |\partial(B_r)_t| dt.$$

Instead of (3.12) we can now use that

$$|\partial(B_r)_t| = |\partial B_r| \left(1 - \frac{t}{r}\right)_+^{d-1}$$

and conclude that for $\sigma \geq 3/2$ and $\Lambda > 0$

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma,d}^{cl} |B_r| \Lambda^{\sigma+d/2} - L_{\sigma,d}^{cl} 2^{-d-2} |\partial B_r| \int_0^1 \left(1 - \frac{s}{4r\sqrt{\Lambda}}\right)_+^{d-1} ds \Lambda^{\sigma+(d-1)/2}.$$

We note that $\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma = 0$ for $\Lambda \leq \lambda_1(B_r)$. For $\Lambda > \lambda_1(B_r)$ we apply the foregoing inequality with $\sigma = 3/2$. Then the method from [AL78] yields that for $\sigma \geq 3/2$ the inequality

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma,d}^{cl} |B_r| \Lambda^{\sigma+d/2} - C_{ba} L_{\sigma,d-1}^{cl} |\partial B_r| \Lambda^{\sigma+(d-1)/2} \quad (3.13)$$

holds with a constant

$$C_{ba} = \frac{j_{d/2-1,1}}{2^{d+1} d \pi^{1/2}} \frac{\Gamma((d+4)/2)}{\Gamma((d+5)/2)} \left(1 - \left(1 - \frac{1}{4j_{d/2-1,1}}\right)^d\right) > 0.$$

4. LOWER BOUNDS ON INDIVIDUAL EIGENVALUES

In order to further estimate the remainder terms, in particular to show that the remainder is negative for all $\Lambda \geq \lambda_1(\Omega)$ - as in (3.13) for the ball - one needs suitable bounds on the ground state $\lambda_1(\Omega)$. We point out the following consequence of the proof of Theorem 4 which might be of independent interest.

Corollary 7. *For any open set $\Omega \subset \mathbb{R}^d$ the estimate*

$$\lambda_1(\Omega) \geq \frac{\pi^2}{l_0^2}$$

holds, where l_0 is given in (3.2).

Proof. Fix $\varepsilon > 0$ and choose a direction $u_0 \in \mathbb{S}^{d-1}$, such that $\sup_{x \in \Omega} l(x, u_0) < l_0 + \varepsilon$. We write $x = (x', t) \in \mathbb{R}^{d-1} \times \mathbb{R}$, where the t -axis is chosen in the direction of u_0 . Let us recall inequality (3.5) from the proof of Theorem 4: For any $\sigma \geq 3/2$

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \mathrm{Tr} V(x', \Lambda)^{\sigma+(d-1)/2} dx',$$

where $V(x', \Lambda)$ denotes the negative part of the operator $-\partial_t^2 - \Lambda$ on $\Omega(x') = \bigcup_{k=1}^{N(x')} J_k(x')$ with Dirichlet boundary conditions at the endpoints of each interval $J_k(x')$. This inequality can be rewritten as

$$\mathrm{Tr}(-\Delta - \Lambda)_-^\sigma \leq L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \sum_{k=1}^{N(x')} \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{|J_k(x')|^2} \right)_+^{\sigma+(d-1)/2} dx'.$$

Our choice of coordinate system implies $|J_k(x')| \leq \sup_{x \in \Omega} l(x, u_0) < l_0 + \varepsilon$ for all $k = 1, \dots, N(x')$ and all $x' \in \mathbb{R}^{d-1}$. It follows that the right hand side of the inequality above is zero for all $\Lambda \leq \pi^2/(l_0 + \varepsilon)^2$. Thus by taking the limit $\varepsilon \rightarrow 0$ we find

$$\sum_{n \in \mathbb{N}} (\Lambda - \lambda_n)_+^\sigma = \mathrm{Tr}(-\Delta - \Lambda)_-^\sigma = 0,$$

for all $\Lambda \leq \pi^2/l_0^2$ and $\lambda_1 \geq \pi^2/l_0^2$ follows. \square

From (3.3) we obtain similar bounds on higher eigenvalues using a method introduced in [Lap97].

Corollary 8. *For any open set $\Omega \subset \mathbb{R}^d$ with finite volume and any $k \in \mathbb{N}$ the estimate*

$$\lambda_k(\Omega) \geq C_d \left(\frac{12}{\pi} \right)^{1/d} \frac{d}{(d+3)^{1+1/d}} \left(\frac{\Gamma((d+3)/2)}{\Gamma(d/2+1)} \right)^{2/d} \frac{k^{2/d}}{|\Omega|^{2/d}} + \frac{1}{l_0^2}$$

holds, with

$$C_d = 4\pi \Gamma(d/2+1)^{2/d}.$$

Proof. Let $N(\Lambda) = \mathrm{Tr}(-\Delta - \Lambda)_-^0$ denote the counting function of the eigenvalues below $\Lambda > 0$. In [Lap97] it is shown that for $\sigma > 0$, and all $\Lambda > 0$, $\tau > 0$

$$N(\Lambda) \leq (\tau\Lambda)^{-\sigma} \mathrm{Tr}(-\Delta - (1+\tau)\Lambda)_-^\sigma. \quad (4.1)$$

If we apply this inequality with $\sigma = 3/2$, we can use (3.3) to estimate

$$N(\Lambda) \leq L_{3/2, d}^{cl} |\Omega| \Lambda^{d/2} \frac{(1+\tau)^{(d+3)/2}}{\tau^{3/2}} \left(1 - \frac{1}{\Lambda(1+\tau)l_0^2} \right)_+^{(d+3)/2}.$$

Minimising the right hand side in $\tau > 0$ yields $\tau_{\min} = 3(\Lambda l_0^2 - 1)/(d\Lambda l_0^2)$ and inserting this we find

$$N(\Lambda) \leq L_{3/2, d}^{cl} |\Omega| \frac{(d+3)^{(d+3)/2}}{3^{3/2} d^{d/2}} \left(\Lambda - \frac{1}{l_0^2} \right)_+^{d/2}.$$

This is equivalent to the claimed result. \square

Remark. Applying the same method to (1.1) with $\sigma = 1$ we recover the Li-Yau inequality (1.5). In the proof of Corollary 8 we have to start from $\sigma = 3/2$, therefore the result is not strong enough to improve (1.5) in general, but one gets improvements for low eigenvalues whenever l_0 is small.

5. FURTHER IMPROVEMENTS IN DIMENSION 2

In this section we further improve Corollary 6 and generalise it to a large class of bounded convex domains $\Omega \subset \mathbb{R}^2$. Here we do not require smoothness, therefore we cannot use (3.12) to estimate inner parallels of the boundary. To find a suitable substitute let w denote the minimal width of Ω and note that for l_0 given in (3.2) the identity

$$w = l_0$$

holds true, see e.g. [BF48]. In the remainder of this section we assume that for all $t > 0$

$$|\partial\Omega_t| \geq \left(1 - \frac{3t}{w}\right)_+ |\partial\Omega|. \quad (5.1)$$

This is true for a large class of convex domains, including the circle, regular polygons and arbitrary triangles. Actually we conjecture that (5.1) holds true for all bounded convex domains in \mathbb{R}^2 .

Furthermore we need a lower bound on the ground state. From Corollary 7 we obtain that for all convex domains $\Omega \subset \mathbb{R}^2$

$$\lambda_1(\Omega) \geq \frac{\pi^2}{w^2} \quad (5.2)$$

holds. One should mention, that the same estimate can be obtained from the inequality $\lambda_1(\Omega) \geq \pi^2/(4r_{in}^2)$, see [Oss77], where r_{in} is the inradius of Ω .

Using similar but more precise methods as in the proof of Theorem 5 we get

Theorem 9. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, convex domain, satisfying (5.1). Then for $\sigma \geq 3/2$ we have*

$$\begin{aligned} \text{Tr}(-\Delta - \Lambda)_-^\sigma &= 0 && \text{if } \Lambda \leq \pi^2/w^2 \text{ and} \\ \text{Tr}(-\Delta - \Lambda)_-^\sigma &\leq L_{\sigma,2}^d |\Omega| \Lambda^{\sigma+1} - C_{co} L_{\sigma,1}^d |\partial\Omega| \Lambda^{\sigma+1/2} && \text{if } \Lambda > \pi^2/w^2, \end{aligned}$$

with a constant

$$C_{co} \geq \frac{11}{9\pi^2} - \frac{3}{20\pi^4} - \frac{2}{5\pi^2} \ln\left(\frac{4\pi}{3}\right) > 0.0642.$$

Proof. The first claim follows directly from (5.2), thus we can assume $\Lambda > \pi^2/w^2$. First we prove the result for $\sigma = 3/2$. Again we can start from (3.7) and we need to estimate $d(x, u)$ in terms of $\delta(x)$, which is just the distance to the boundary, since Ω is convex.

Fix $x \in \Omega$. Since Ω is convex and smooth we can choose $u_0 \in \mathbb{S}^{d-1}$, such that $d(x, u_0) = \delta(x)$. We can assume $u_0 = (1, 0, \dots, 0)$ and put $\mathbb{S}_+^{d-1} = \{u \in \mathbb{R}^d : u_1 > 0\}$.

Let a be the intersection point of the semi-axes $\{x + tu_0, t > 0\}$ with $\partial\Omega$ and for arbitrary $u \in \mathbb{S}_+^{d-1}$ let b_u be the intersection point of $\{x + tu, t > 0\}$ with the plane through a , orthogonal to u_0 . We find $d(x, u) \leq |x - b_u|$ and if θ_u denotes the angle between u_0 and u , we find

$$d(x, u) \leq |x - b_u| = \frac{|x - a|}{\cos \theta_u} = \frac{\delta(x)}{\cos \theta_u}.$$

Using (3.7) and taking into account that $d(x, u) = d(x, -u)$ we can estimate

$$\begin{aligned} \text{Tr}(-\Delta - \Lambda)_-^{3/2} &\leq L_{3/2,2}^d \Lambda^{5/2} \int_{\Omega} \frac{2}{\pi} \int_{\theta_0}^{\pi/2} \left(1 - \frac{\cos^2(\theta)}{4\Lambda\delta(x)^2}\right)^{5/2} d\theta dx \\ &\leq \frac{\Lambda^{5/2}}{5\pi^2} \int_{\Omega} \int_{\theta_0}^{\pi/2} \left(1 - \frac{\cos^2(\theta)}{2\Lambda\delta(x)^2} + \frac{\cos^4(\theta)}{16\Lambda^2\delta(x)^4}\right) d\theta dx, \end{aligned} \quad (5.3)$$

where $\theta_0 = 0$ if $\delta(x) \geq 1/(2\sqrt{\Lambda})$ and $\theta_0 = \arccos(2\delta(x)\sqrt{\Lambda})$ if $\delta(x) < 1/(2\sqrt{\Lambda})$. We set $u_0 = \min(1, 2\delta(x)\sqrt{\Lambda})$ and calculate

$$\begin{aligned} & \int_{\theta_0}^{\pi/2} \left(1 - \frac{\cos^2(\theta)}{2\Lambda\delta(x)^2} + \frac{\cos^4(\theta)}{16\Lambda^2\delta(x)^4} \right) d\theta \\ &= \frac{\pi}{2} - \theta_0 - \frac{\arcsin(u_0) - u_0\sqrt{1-u_0^2}}{4\Lambda\delta(x)^2} + \frac{3\arcsin(u_0) - u_0(2u_0^2+3)\sqrt{1-u_0^2}}{128\Lambda^2\delta(x)^4}. \end{aligned}$$

Inserting this back into (5.3) yields

$$\mathrm{Tr}(-\Delta - \Lambda)_-^{3/2} \leq \frac{1}{10\pi} |\Omega| \Lambda^{5/2} - \Lambda^{5/2} (I_1 + I_2), \quad (5.4)$$

where

$$I_1 = \int_{\{\delta(x) \geq 1/(2\sqrt{\Lambda})\}} \frac{1}{10\pi} \left(\frac{1}{4\Lambda\delta(x)^2} - \frac{3}{128\Lambda^2\delta(x)^4} \right) dx$$

and

$$\begin{aligned} I_2 = & \int_{\{\delta(x) < 1/(2\sqrt{\Lambda})\}} \frac{1}{5\pi^2} \left(\arccos(2\sqrt{\Lambda}\delta(x)) + \frac{\arcsin(2\sqrt{\Lambda}\delta(x))}{4\Lambda\delta(x)^2} - \frac{\sqrt{1-4\Lambda\delta(x)^2}}{2\sqrt{\Lambda}\delta(x)} \right. \\ & \left. - \frac{3\arcsin(2\sqrt{\Lambda}\delta(x))}{128\Lambda^2\delta(x)^4} + \frac{(8\Lambda\delta(x)^2+3)\sqrt{1-4\Lambda\delta(x)^2}}{64\Lambda^{3/2}\delta(x)^3} \right) dx. \end{aligned}$$

First we turn to

$$I_1 = \frac{1}{10\pi} \int_{1/(2\sqrt{\Lambda})}^{\infty} |\partial\Omega_t| \left(\frac{1}{4\Lambda t^2} - \frac{3}{128\Lambda^2 t^4} \right) dt.$$

Note that the term in brackets is positive, thus after substituting $s = 2\sqrt{\Lambda}t$ we can use (5.1) and $\Lambda > \pi^2/w^2$ to obtain

$$\begin{aligned} I_1 &\geq \frac{1}{20\pi} \frac{|\partial\Omega|}{\sqrt{\Lambda}} \int_1^{\infty} \left(1 - \frac{3s}{2\pi} \right)_+ \left(\frac{1}{s^2} - \frac{3}{8s^4} \right) ds \\ &= \frac{1}{20\pi} \frac{|\partial\Omega|}{\sqrt{\Lambda}} \left(\frac{7}{8} - \frac{39}{32\pi} - \frac{27}{128\pi^3} - \frac{3}{2\pi} \ln\left(\frac{2\pi}{3}\right) \right). \end{aligned}$$

Similarly we can treat I_2 and get

$$I_2 \geq \frac{1}{10\pi^2} \frac{|\partial\Omega|}{\sqrt{\Lambda}} \left(\frac{557}{192} - \frac{7\pi}{16} - \frac{3}{2\pi} \int_0^1 \frac{\arcsin(s)}{s} ds \right).$$

In view of $L_{3/2,1}^c = 3/16$ we can write

$$I_1 + I_2 \geq L_{3/2,1}^c \frac{|\partial\Omega|}{\sqrt{\Lambda}} \left(\frac{11}{9\pi^2} - \frac{3}{20\pi^4} - \frac{2}{5\pi^2} \ln\left(\frac{2\pi}{3}\right) - \frac{2}{5\pi^2} \ln(2) \right)$$

and inserting this into (5.4) yields the claim in the case $\sigma = 3/2$.

To prove the estimate for $\sigma > 3/2$ we again refer to [AL78] and use the identity

$$\mathrm{Tr}(-\Delta - \Lambda)_-^{\sigma} = \frac{1}{B(\sigma - 3/2, 5/2)} \int_0^{\infty} \tau^{\sigma-5/2} \mathrm{Tr}(-\Delta - (\Lambda - \tau))_-^{3/2} d\tau,$$

from which the general result follows. \square

Now we can apply the same arguments that lead to Corollary 8 to derive lower bounds on individual eigenvalues.

Corollary 10. *Let $\Omega \subset \mathbb{R}^2$ be a bounded, convex domain, satisfying (5.1). Then for $k \in \mathbb{N}$ and any $\alpha \in (0, 1)$ the estimate*

$$\frac{\lambda_k(\Omega)}{1-\alpha} \geq 10\pi\alpha^{3/2} \frac{k}{|\Omega|} + \frac{15\pi C_{co}}{8} \frac{|\partial\Omega|}{|\Omega|} \sqrt{10\pi\alpha^{3/2} \frac{k}{|\Omega|} + \frac{225\pi^2 C_{co}^2}{256} \frac{|\partial\Omega|^2}{|\Omega|^2}} + \frac{225\pi^2 C_{co}^2}{128} \frac{|\partial\Omega|^2}{|\Omega|^2}$$

holds, with the constant C_{co} given in Theorem 9.

Proof. Applying (4.1) and Theorem 9 with $\sigma = 3/2$ yields

$$N(\Lambda) \leq L_{3/2,2}^{cl} |\Omega| \Lambda \frac{(1+\tau)^{5/2}}{\tau^{3/2}} - C_{co} L_{3/2,1}^{cl} |\partial\Omega| \sqrt{\Lambda} \frac{(1+\tau)^2}{\tau^{3/2}}$$

for any $\tau > 0$ and $\Lambda > \pi^2/w^2$. With $\tau = \alpha/(1-\alpha)$, $\alpha \in (0, 1)$, this is equivalent to the claimed estimate. \square

Remark. Given a fixed ratio $|\partial\Omega|/|\Omega|$ one can optimise the foregoing estimate in $\alpha \in (0, 1)$, depending on $k \in \mathbb{N}$. As mentioned in the remark after Corollary 8, the result cannot improve the Li-Yau inequality (1.5) in general, since we have to apply (4.1) with $\sigma = 3/2$ instead of $\sigma = 1$. However, the estimates obtained from Corollary 10 are stronger than (1.5) for low eigenvalues and the improvements depend on the ratio $|\partial\Omega|/|\Omega|$.

In particular, one can use the isoperimetric inequality, namely that $|\partial\Omega| \geq 2(\pi|\Omega|)^{1/2}$ for all $\Omega \subset \mathbb{R}^2$, to derive general improvements of the Li-Yau inequality (1.5) for low eigenvalues. Indeed, from (1.5) we get

$$\lambda_k(\Omega) \geq \frac{2\pi k}{|\Omega|},$$

while optimising the estimate of Corollary 10 with $|\partial\Omega| = 2(\pi|\Omega|)^{1/2}$, we find that for any convex domain, satisfying (5.1),

$$\lambda_2(\Omega) > \frac{15.03}{|\Omega|} > \frac{4\pi}{|\Omega|}, \quad \lambda_3(\Omega) > \frac{21.52}{|\Omega|} > \frac{6\pi}{|\Omega|}, \quad \dots, \quad \lambda_{23}(\Omega) > \frac{144.58}{|\Omega|} > \frac{46\pi}{|\Omega|}.$$

In this way we can improve (1.5) in convex domains for all eigenvalues $\lambda_k(\Omega)$ with $k \leq 23$.

Finally let us make a remark about the square $Q_l = (0, l) \times (0, l) \subset \mathbb{R}^2$, $l > 0$. Using the methods introduced in section 2 one can establish the following two-dimensional version of Lemma 1: Choose a coordinate system $(x_1, x_2) \in \mathbb{R}^2$ with axes parallel to the sides of the square and for $x \in Q_l$ put

$$\delta(x_i) = \min(x_i, l - x_i), \quad i = 1, 2.$$

Then for $\sigma \geq 1$ and all $\Lambda > 0$ the estimate

$$\begin{aligned} \text{Tr}(-\Delta - \Lambda)_-^\sigma &= \sum_{m,n \in \mathbb{N}} \left(\Lambda - \frac{\pi^2}{l^2} (n^2 + m^2) \right)_+^\sigma \\ &\leq L_{\sigma,2}^{cl} \int_0^l \int_0^l \left(\Lambda - C_{sq} \left(\frac{1}{\delta(x_1)} + \frac{1}{\delta(x_2)} \right)^2 \right)_+^{\sigma+1} dx_1 dx_2 \end{aligned}$$

holds with a constant $C_{sq} > 1/10$.

APPENDIX A. PROOF OF LEMMA 2 AND LEMMA 3

A.1. Proof of Lemma 2. For $A \in \mathbb{R}$ let \bar{A} and \tilde{A} denote the integer and fractional part of A respectively. Then we can calculate

$$\begin{aligned} \sum_k \left(1 - \frac{k^2}{A^2}\right)_+ &= \sum_{k=1}^{\bar{A}} \left(1 - \frac{k^2}{A^2}\right) = \bar{A} - \frac{1}{A^2} \left(\frac{\bar{A}^3}{3} + \frac{\bar{A}^2}{2} + \frac{\bar{A}}{6}\right) \\ &= \frac{2A}{3} - \frac{1}{2} - \frac{1}{6A} + \tilde{A}(1 - \tilde{A})\frac{1}{A} + \tilde{A} \left(1 - 3\tilde{A} + 2\tilde{A}^2\right) \frac{1}{6A^2}. \end{aligned}$$

From $0 \leq \tilde{A} < 1$ we conclude $\tilde{A}(1 - \tilde{A}) \leq 1/4$ and $\tilde{A} \left(1 - 3\tilde{A} + 2\tilde{A}^2\right) \leq \sqrt{3}/18$ and we get

$$\sum_k \left(1 - \frac{k^2}{A^2}\right)_+ \leq \frac{2A}{3} - \frac{1}{2} + \frac{1}{12A} + \frac{\sqrt{3}}{108A^2}. \quad (\text{A.1})$$

To estimate the right hand side of (2.1) note that

$$\int \left(1 - \frac{1}{s^2}\right)^{3/2} ds = \left(1 + \frac{1}{2s^2}\right) \sqrt{s^2 - 1} + \frac{3}{2} \arctan\left(\frac{1}{\sqrt{s^2 - 1}}\right),$$

thus

$$\frac{2}{3\pi} \int_1^{\pi A} \left(1 - \frac{1}{s^2}\right)^{3/2} ds = \frac{2\pi^2 A^2 + 1}{3\pi^2 A} \frac{\sqrt{\pi^2 A^2 - 1}}{\pi A} + \frac{1}{\pi} \arctan\left(\frac{1}{\sqrt{\pi^2 A^2 - 1}}\right) - \frac{1}{2}.$$

Now we can insert the elementary estimates

$$\arctan\left(\frac{1}{\sqrt{\pi^2 A^2 - 1}}\right) \geq \frac{1}{\pi A} \quad \text{and} \quad \frac{\sqrt{\pi^2 A^2 - 1}}{\pi A} \geq 1 - \frac{1}{2\pi^2 A^2} - \frac{a}{A^4}$$

both valid for $A \geq 2$, where we write $a = 16 - 2/\pi^2 - 8\sqrt{4\pi^2 - 1}/\pi$ for simplicity. We get

$$\frac{2}{3\pi} \int_1^{\pi A} \left(1 - \frac{1}{s^2}\right)^{3/2} ds \geq \frac{2A}{3} - \frac{1}{2} + \frac{1}{\pi^2 A} - \frac{1}{6\pi^4 A^3} - \frac{a}{3} \left(\frac{2}{A^3} + \frac{1}{\pi^2 A^5}\right) \quad (\text{A.2})$$

for all $A \geq 2$. From (A.1) and (A.2) we deduce that (2.1) holds true for all $A \geq 2$, since

$$\left(\frac{1}{\pi^2} - \frac{1}{12}\right) A^4 - \frac{\sqrt{3}}{108} A^3 - \left(\frac{1}{6\pi^4} + \frac{2a}{3}\right) A^2 - \frac{a}{3\pi^2} \geq 0$$

for all $A \geq 2$.

Note that (2.1) is trivial for $1/\pi \leq A \leq 1$, since the left hand side equals zero. The remaining case $1 \leq A \leq 2$ can be checked by hand.

A.2. Proof of Lemma 3. We assume $I = (0, l)$, substitute $t = s\sqrt{c/\Lambda}$ and write

$$\int_0^l \left(\Lambda - \frac{c}{\delta(t)^2}\right)_+^{\sigma+1/2} dt = 2\sqrt{c}\Lambda^\sigma \int_1^{l\sqrt{\Lambda}/(2\sqrt{c})} \left(1 - \frac{1}{s^2}\right)^{\sigma+1/2} ds.$$

The claim of the Lemma follows, if we show that

$$2\sqrt{c}L_{\sigma,1}^{cl} \int_1^{l\sqrt{\Lambda}/(2\sqrt{c})} \left(1 - \frac{1}{s^2}\right)^{\sigma+1/2} ds - \sum_k \left(1 - \frac{\pi^2 k^2}{\Lambda l^2}\right)_+^\sigma = \frac{1}{2} - \sqrt{c} + o(1)$$

as $\Lambda \rightarrow \infty$. With $A = l\sqrt{\Lambda}/\pi$ this is equivalent to

$$2\sqrt{c} L_{\sigma,1}^{cl} \int_1^{\pi A/(2\sqrt{c})} \left(1 - \frac{1}{s^2}\right)^{\sigma+1/2} ds - \sum_k \left(1 - \frac{k^2}{A^2}\right)_+^{\sigma} = \frac{1}{2} - \sqrt{c} + o(1) \quad (\text{A.3})$$

as $A \rightarrow \infty$.

It is easy to see that

$$\sum_k \left(1 - \frac{k^2}{A^2}\right)_+^{\sigma} = \frac{A}{2} B\left(\sigma + 1, \frac{1}{2}\right) - \frac{1}{2} + o(1)$$

as $A \rightarrow \infty$. Moreover, we claim

$$\int_1^{\pi A/(2\sqrt{c})} \left(1 - \frac{1}{s^2}\right)^{\sigma+1/2} ds = \frac{\pi A}{2\sqrt{c}} + \frac{1}{2} B\left(-\frac{1}{2}, \sigma + \frac{3}{2}\right) + o(1) \quad (\text{A.4})$$

as $A \rightarrow \infty$ and (A.3) follows from $L_{\sigma,1}^{cl} = B(\sigma + 1, 1/2)/(2\pi)$, if we can establish (A.4). Let us write

$$\left(1 - \frac{1}{s^2}\right)^m = \sum_{k \geq 0} (-1)^k \binom{m}{k} s^{-2k} = \sum_{k \geq 0} \binom{k - m - 1}{k} s^{-2k}$$

for $m \geq 1$ and note that the sum is finite if $m \in \mathbb{N}$, while the sum converges uniformly on $s \in [1, \infty)$ if $m \notin \mathbb{N}$. Hence we have

$$\int_1^y \left(1 - \frac{1}{s^2}\right)^m ds = y + \sum_{k \geq 0} \binom{k - m - 1}{k} \frac{1}{2k - 1} + o(1)$$

as $y \rightarrow \infty$. Using that

$$\sum_{k \geq 0} \binom{k - m - 1}{k} \frac{1}{2k - 1} = \frac{1}{2} B\left(-\frac{1}{2}, m + 1\right)$$

we obtain

$$\int_1^y \left(1 - \frac{1}{s^2}\right)^m ds = y + \frac{1}{2} B\left(-\frac{1}{2}, m + 1\right) + o(1)$$

as $y \rightarrow \infty$, which is equivalent to (A.4). This finishes the proof of Lemma 3.

REFERENCES

- [AL78] M. Aizenman and E.H. Lieb, *On semi-classical bounds for eigenvalues of Schrödinger operators*, Phys. Lett. **66** (1978), 427–429.
- [Ber72] F. A. Berezin, *Covariant and contravariant symbols of operators*, Izv. Akad. Nauk SSSR Ser. Mat. **13** (1972), 1134–1167.
- [BF48] T. Bonnesen and W. Fenchel, *Theorie der konvexen Körper*, Chelsea Publishing Company, New York, 1948.
- [CH24] R. Courant and D. Hilbert, *Methoden der mathematischen Physik*, Springer, Berlin, 1924.
- [Dav94] E. B. Davies, *Spectral theory and differential operators*, Cambridge Studies in Advances Mathematics, vol. 42, Cambridge University Press, Cambridge, 1994.
- [Dav99] ———, *A review of Hardy inequalities*, The Maz'ya anniversary collection, Vol. 2, Oper. Theory Adv. Appl., vol. 110, Birkhauser, Basel, 1999, pp. 55–67.
- [EF06] T. Ekholm and R. L. Frank, *On Lieb-Thirring inequalities for Schrödinger operators with virtual level*, Comm. Math. Phys. **264** (2006), 725–740.
- [ELW04] P. Exner, H. Linde, and T. Weidl, *Lieb-Thirring inequalities for geometrically induced bound states*, Lett. Math. Phys. **70** (2004), no. 1, 83–95.

- [FL08] R. L. Frank and A. Laptev, *Spectral inequalities for Schrödinger operators with surface potentials*, Spectral theory of differential operators, Amer. Math. Soc. Transl. Ser. 2, vol. 225, Amer. Math. Soc., Providence, RI, 2008, pp. 91–102.
- [FLS08] R. L. Frank, E. H. Lieb, and R. Seiringer, *Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators*, J. Amer. Math. Soc. **21** (2008), no. 4, 925–950.
- [FLU02] J. K. Freericks, E. H. Lieb, and D. Ueltschi, *Segregation in the Falicov-Kimball model*, Comm. Math. Phys. **227** (2002), no. 2, 243–279.
- [FLV95] J. Fleckinger, M. Levitin, and D. Vassiliev, *Heat equation on the triadic von Koch snowflake: Asymptotic and numerical analysis*, Proc. London Math. Soc. **71** (1995), no. 3, 372–396.
- [Fra09] R. L. Frank, *A simple proof of Hardy-Lieb-Thirring inequalities*, Comm. Math. Phys. **290** (2009), no. 2, 789–800.
- [FV93] J. Fleckinger and D. G. Vassiliev, *An example of a two-term asymptotics for the “counting function” of a fractal drum*, Trans. Amer. Math. Soc. **337** (1993), no. 1, 99–116.
- [Gug77] H. W. Guggenheimer, *Differential geometry*, New York, Dover, 1977.
- [GW10] L. Geisinger and T. Weidl, *Universal bounds for traces of the Dirichlet Laplace operator*, J. Lond. Math. Soc. doi 10.1112/jlms/jdq033 (2010).
- [HHL02] M. HoffmannOstenhof, T. HoffmannOstenhof, and A. Laptev, *A geometrical version of Hardy’s inequality*, J. Funct. Anal. **189** (2002), 539–548.
- [Hör68] L. Hörmander, *The spectral function of an elliptic operator*, Acta Math. **121** (1968), 193–218.
- [Ily09] A. A. Ilyin, *Lower bounds for the spectrum of the Laplace and Stokes operators*, arXiv.org: 0909.2181v1 (2009).
- [Ivr80] V. Ya. Ivrii, *On the second term of the spectral asymptotics for the Laplace-Beltrami operator on manifolds with boundary*, Funtsional. Anal. i Prilozhen. **14** (1980), no. 2, 25–34.
- [Ivr98] ———, *Microlocal analysis and precise spectral asymptotics*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998.
- [KVVW09] H. Kovařík, S. Vugalter, and T. Weidl, *Two dimensional Berezin-Li-Yau inequalities with a correction term*, Comm. Math. Phys. **287** (2009), no. 3, 959–981.
- [Lap91] M. L. Lapidus, *Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl-Berry conjecture*, Trans. Amer. Math. Soc. **325** (1991), no. 2, 465–529.
- [Lap97] A. Laptev, *Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces*, J. Funct. Anal. **151** (1997), no. 2, 531–545.
- [LW00] A. Laptev and T. Weidl, *Sharp Lieb-Thirring inequalities in high dimensions*, Acta Math. **184** (2000), no. 1, 87–111.
- [LY83] P. Li and S. T. Yau, *On the Schrödinger equation and the eigenvalue problem*, Comm. Math. Phys. **88** (1983), no. 3, 309–318.
- [Mel80] R. B. Melrose, *Weyl’s conjecture for manifolds with concave boundary*, Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979). Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., 1980, pp. 257–274.
- [Mel03] A.D. Melás, *A lower bound for sums of eigenvalues of the Laplacian*, Amer. Math. Soc. **131** (2003), 631–636.
- [Oss77] R. Osserman, *A note on Hayman’s theorem on the bass note of a drum*, Comment. Math. Helv. **52** (1977), no. 4, 545–555.
- [SV97] Y. Safarov and D. Vassiliev, *The asymptotic distribution of eigenvalues of partial differential operators*, Translations of Mathematical Monographs, 155, American Mathematical Society, Providence, RI, 1997.
- [vdB84] M. van den Berg, *A uniform bound on trace ($e^{t\Delta}$) for convex regions in \mathbb{R}^n with smooth boundaries*, Comm. Math. Phys. **92** (1984), no. 4, 525–530.
- [Wei08] T. Weidl, *Improved Berezin-Li-Yau inequalities with a remainder term*, Amer. Math. Soc. Transl. **225** (2008), no. 2, 253–263.
- [Wey12] H. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)*, Math. Ann. **71** (1912), no. 4, 441–479.
- [YB61] I. M. Yaglom and V. G. Boltyanski, *Convex figures*, Holt, Reinhart and Winston, New York, 1961.
- [Yol09] S. Y. Yolcu, *An improvement to a Berezin-Li-Yau type inequality for the Klein-Gordon operator*, arXiv.org: 0909.4132v1 (2009).

Leander Geisinger
Universität Stuttgart
Pfaffenwaldring 57
D - 70569 Stuttgart
E-Mail: geisinger@mathematik.uni-stuttgart.de

Ari Laptev
Imperial College London
180 Queen's Gate
London SW7 2AZ
UK
E-Mail: a.laptev@imperial.ac.uk

Timo Weidl
Universität Stuttgart
Pfaffenwaldring 57
D - 70569 Stuttgart
E-Mail: weidl@mathematik.uni-stuttgart.de

Erschienenene Preprints ab Nummer 2007/001

Komplette Liste: <http://www.mathematik.uni-stuttgart.de/preprints>

- 2010/009 *Geisinger, L.; Laptev, A.; Weidl, T.:* Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010/008 *Poppitz, S.; Stroppel, M.:* Polarities of Schellhammer Planes
- 2010/007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.:* Non-existence of isomorphisms between certain unitals
- 2010/006 *Höllig, K.; Hörner, J.; Hoffacker, A.:* Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods
- 2010/005 *Kaltenbacher, B.; Walk, H.:* On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010/004 *Kühnel, W.; Solanes, G.:* Tight surfaces with boundary
- 2010/003 *Kohler, M.; Walk, H.:* On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010/002 *Gulde, M.; Stroppel, M.:* Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010/001 *Leitner, F.:* Examples of almost Einstein structures on products and in cohomogeneity one
- 2009/008 *Griesemer, M.; Zenk, H.:* On the atomic photoeffect in non-relativistic QED
- 2009/007 *Griesemer, M.; Moeller, J.S.:* Bounds on the minimal energy of translation invariant n-polaron systems
- 2009/006 *Demirel, S.; Harrell II, E.M.:* On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009/005 *Bächle, A.; Kimmerle, W.:* Torsion subgroups in integral group rings of finite groups
- 2009/004 *Geisinger, L.; Weidl, T.:* Universal bounds for traces of the Dirichlet Laplace operator
- 2009/003 *Walk, H.:* Strong laws of large numbers and nonparametric estimation
- 2009/002 *Leitner, F.:* The collapsing sphere product of Poincaré-Einstein spaces
- 2009/001 *Brehm, U.; Kühnel, W.:* Lattice triangulations of E^3 and of the 3-torus
- 2008/006 *Kohler, M.; Krzyżak, A.; Walk, H.:* Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps
- 2008/005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.:* Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems
- 2008/004 *Leitner, F.:* Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008/003 *Effenberger, F.; Kühnel, W.:* Hamiltonian submanifolds of regular polytope
- 2008/002 *Hertweck, M.; Hofert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups $PSL(2, q)$
- 2008/001 *Kovarik, H.; Vugalter, S.; Weidl, T.:* Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007/006 *Weidl, T.:* Improved Berezin-Li-Yau inequalities with a remainder term
- 2007/005 *Frank, R.L.; Loss, M.; Weidl, T.:* Polya's conjecture in the presence of a constant magnetic field
- 2007/004 *Ekholm, T.; Frank, R.L.; Kovarik, H.:* Eigenvalue estimates for Schrödinger operators on metric trees
- 2007/003 *Lesky, P.H.; Racke, R.:* Elastic and electro-magnetic waves in infinite waveguides
- 2007/002 *Teufel, E.:* Spherical transforms and Radon transforms in Moebius geometry
- 2007/001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions