

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Sharp spectral estimates in domains of infinite volume

Leander Geisinger, Timo Weidl

Preprint 2011/001

**Universität
Stuttgart**

**Fachbereich
Mathematik**

Sharp spectral estimates in domains of infinite volume

Leander Geisinger, Timo Weidl

Preprint 2011/001

Fachbereich Mathematik
Fakultät Mathematik und Physik
Universität Stuttgart
Pfaffenwaldring 57
D-70 569 Stuttgart

E-Mail: preprints@mathematik.uni-stuttgart.de

WWW: <http://www.mathematik.uni-stuttgart.de/preprints>

ISSN **1613-8309**

© Alle Rechte vorbehalten. Nachdruck nur mit Genehmigung des Autors.

LaTeX-Style: Winfried Geis, Thomas Merkle

1. INTRODUCTION

Let $V(x)$ be a non-negative function on an open set $\Omega \subset \mathbb{R}^d$, $d \geq 1$. In this article we study the negative spectrum of Schrödinger operators

$$H_\Omega = -\Delta - V$$

defined in $L^2(\Omega)$ with Dirichlet conditions on the boundary of Ω . More precisely, one defines H_Ω to be the self-adjoint operator generated by the quadratic form

$$\langle u, H_\Omega u \rangle = \int_\Omega |\nabla u(x)|^2 dx - \int_\Omega V(x) |u(x)|^2 dx,$$

with form domain $H_0^1(\Omega)$, see [BS87] for details. We always assume that H_Ω has purely discrete spectrum. Then the negative spectrum of H_Ω consists of finitely many eigenvalues $-\lambda_1 \leq -\lambda_2 \leq \dots - \lambda_N < 0$, $N < \infty$, counted with multiplicity. In general, these eigenvalues cannot be calculated explicitly and for large N it is difficult to approximate them numerically. Hence, to deduce information about the eigenvalues one studies also the Riesz means

$$R_\sigma(V; \Omega) = \text{Tr}(H_\Omega)_-^\sigma = \sum_{k=1}^N \lambda_k^\sigma$$

of order $\sigma \geq 0$ and their dependence on Ω and V .

The first rigorous step in this direction dates back to H. Weyl, R. Courant and D. Hilbert [Wey12, CH24] who calculated the semiclassical limit of the eigenvalues in the case of a constant potential. To state the general result let us introduce a scaling parameter $\lambda > 0$ and replace the potential V by λV . Then for $\sigma \geq 0$ and $V \in L^{\sigma+d/2}(\Omega)$ the limit

$$R_\sigma(\lambda V; \Omega) = L_{\sigma,d}^{cl} \int_\Omega V(x) dx \lambda^{\sigma+d/2} + o(\lambda^{\sigma+d/2}), \quad \lambda \rightarrow \infty, \quad (1)$$

holds with the semiclassical constant

$$L_{\sigma,d}^{cl} = \frac{\Gamma(\sigma+1)}{(4\pi)^{d/2} \Gamma(\sigma + \frac{d}{2} + 1)},$$

see e.g. [RS78]. To get information about finite potentials one needs to supplement this asymptotic result with uniform estimates. In [LT76] it was shown that for $\Omega = \mathbb{R}^d$ and $\sigma > \max\{0, 1 - d/2\}$ the estimate

$$R_\sigma(V; \mathbb{R}^d) \leq L_{\sigma,d} \int_{\mathbb{R}^d} V(x)^{\sigma+d/2} dx$$

holds with certain positive constants $L_{\sigma,d}$. These inequalities have many important applications, for example, in proving the stability of matter [Lie97, LS10].

Finding the best constants for which the Lieb-Thirring inequalities hold, poses a substantial mathematical challenge. In [LW00] the inequalities were established for $\sigma \geq 3/2$ with the sharp constants $L_{\sigma,d} = L_{\sigma,d}^{cl}$. This result immediately implies that for any open set $\Omega \subset \mathbb{R}^d$, $\sigma \geq 3/2$, and any non-negative potential $V \in L^{\sigma+d/2}(\Omega)$

$$R_\sigma(V; \Omega) \leq L_{\sigma,d}^{cl} \int_\Omega V(x)^{\sigma+d/2} dx. \quad (2)$$

If $V \in L^{\sigma+d/2}(\Omega)$ then both (1) and (2) hold and we see that the bound (2) is sharp: It shows the correct power of V and holds with the sharp constant.

In this article we are interested in the case $V \notin L^{\sigma+d/2}(\Omega)$, where the bound (2) and even the asymptotics (1) must fail and one needs to find a new approach to get sharp uniform bounds on eigenvalues means. If $V \notin L^{\sigma+d/2}(\Omega)$ the leading order of the semiclassical limit depends on the potential V and on the geometry of Ω and it is challenging to find estimates that take these dependencies into account.

Let us discuss the case of constant potential $V \equiv \Lambda > 0$ on Ω in more detail. If Ω is bounded then the semiclassical limit (1) reads as

$$R_\sigma(\Lambda; \Omega) = L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2} + o(\Lambda^{\sigma+d/2}), \quad \sigma \geq 0, \quad \Lambda \rightarrow \infty, \quad (3)$$

where $|\Omega|$ denotes the volume of Ω . In this case the asymptotic results are supplemented by the Berezin-Lieb-Li-Yau inequality [Ber72, Lie73, LY83]: For $\sigma \geq 1$

$$R_\sigma(\Lambda; \Omega) \leq L_{\sigma,d}^{cl} |\Omega| \Lambda^{\sigma+d/2}, \quad \Lambda > 0. \quad (4)$$

Again, the constant in this inequality is sharp and cannot be improved. However, under certain conditions on the geometry of Ω a negative second term exists in the semiclassical limit (3), see [Ivr80, Hör85, SV97, Ivr98, FG10], and the question arises whether (4) can be improved by an additional negative correction term. Recently, several results have been found giving answer to this question [FLU02, Mel03, Wei08, KVV09, GW10, GLW11]. In [FLU02] the corresponding sharp estimate for the discrete Laplacian was improved by a negative remainder term capturing the properties of the second term of the semiclassical limit. The first uniform improvement for the continuous Laplacian is due to A. Melás [Mel03]. He improved the estimate (4) for $\sigma \geq 1$, however, the remainder does not reflect the correct order of the second term of the semiclassical limit.

In [Wei08] this was improved in the case $\sigma \geq 3/2$. Using an inductive argument based on operator-valued Lieb-Thirring inequalities [LW00] the Berezin inequality (4) was strengthened by a negative remainder term of correct order in comparison with the second term of the semiclassical limit. Here we are not concerned with the remainder term but we apply the same inductive argument to derive sharp spectral inequalities in domains of infinite volume.

However, for unbounded domains Ω even the discreteness of the spectrum of the Dirichlet Laplacian is no longer guaranteed. A necessary condition is the so called quasi-boundedness of Ω [AF03] which is satisfied, by definition, if

$$\lim_{\substack{x \in \Omega \\ |x| \rightarrow \infty}} \text{dist}(x, \partial\Omega) = 0.$$

But even for quasi-bounded domains (3) and (4) must fail if the volume of Ω is infinite. In this article we show that one can nevertheless prove uniform bounds on the eigenvalue means for certain domains with infinite volume. In this case the leading order of the semiclassical limit depends on the geometry of Ω , see e.g. [Fle78, Sim83]. However, applying the induction-in-the-dimension argument from [Wei08] we can prove sharp estimates valid for all $\Lambda > 0$ that capture the correct asymptotic behavior.

If the potential V is not constant the situation is more difficult. The same inductive argument can still be used to reduce the problem to one dimension. But in contrast to the case of constant potential the eigenvalues of the resulting one-dimensional operator cannot be calculated explicitly. Therefore we have to study the one-dimensional problem in more detail. In particular, we have to analyze the effect of different boundary conditions on the eigenvalues. The result yields an improved version of the semiclassical bound (2). Again, this

sharp Lieb-Thirring inequality with remainder term can be applied in situations, where all known results – in particular (1) and (2) – fail.

The remainder of the article is structured as follows. First we mention some key ingredients of the proofs. In particular, we review the induction-in-the-dimension argument from [Wei08] and adapt it to our needs here. This is done in Section 2.

In Section 3 we consider constant potentials on domains with infinite volume. We give examples, where the leading order of the semiclassical limit depends on the geometry of the domain Ω . In this situation we derive sharp upper bounds on the eigenvalue means.

The last part of the article is devoted to the general setting of non-constant potentials. In Section 4 we first analyze the effect of different boundary conditions on the eigenvalues of one-dimensional Schrödinger operators. We find an improvement of (2) that can be generalized to higher dimensions. Finally, we give an example to show that the result applies for certain potentials $V \notin L^{\sigma+d/2}(\Omega)$.

2. INDUCTION IN THE DIMENSION

In this section we prove an inequality reducing estimates for eigenvalue means of the operator H_Ω to estimates for one-dimensional Schrödinger operators. The proof relies on a lifting technique from [Lap97] and uses operator-valued Lieb-Thirring inequalities [LW00]. Here we follow the proof from [Wei08], where this approach of induction-in-the-dimension is employed to derive improvements of (4) for constant potentials.

Fix a Cartesian coordinate system in \mathbb{R}^d and for $x \in \mathbb{R}^d$ write $x = (x', t) \in \mathbb{R}^{d-1} \times \mathbb{R}$. For $x' \in \mathbb{R}^{d-1}$ consider one-dimensional sections $\Omega(x') = \{t \in \mathbb{R} : (x', t) \in \Omega\}$. If not empty, each section $\Omega(x')$ consists of at most countably many open intervals $J_k(x') \subset \mathbb{R}$, $k = 1, \dots, N(x') \leq \infty$.

For $x = (x', t) \in \Omega$ put $V_{x'}(t) = V(x)$ and let the one-dimensional Schrödinger operators

$$H_k(x') = -\frac{d^2}{dt^2} - V_{x'}, \quad k = 1, \dots, N(x'),$$

be defined in $L^2(J_k(x'))$ with Dirichlet boundary conditions. Finally let

$$W(x', V) = \bigoplus_{k=1}^{N(x')} H_k(x')_- \tag{5}$$

be the negative part of the Schrödinger operator $-d^2/dt^2 - V_{x'}$ given on $\Omega(x')$ with Dirichlet boundary conditions at the endpoints of each interval forming $\Omega(x')$, that is, on the boundary of $\Omega(x')$.

Using operator-valued Lieb-Thirring inequalities one can estimate eigenvalue means of H_Ω in terms of $W(x', V)$.

Proposition 2.1. *For $\sigma \geq 3/2$ we have*

$$R_\sigma(V; \Omega) \leq L_{\sigma, d-1}^{\text{cl}} \int_{\mathbb{R}^{d-1}} \text{Tr} W(x', V)^{\sigma+(d-1)/2} dx'.$$

Remark. In the case of constant potential $V \equiv \Lambda > 0$ the trace of $W(x', \Lambda)$ can be evaluated explicitly. If Ω is bounded, a detailed analysis of the resulting estimate leads to improved Berezin-Li-Yau inequalities with a remainder term capturing the properties of the second term of the semiclassical limit [Wei08, GW10, GLW11].

Proof of Proposition 2.1. We consider the quadratic form $\langle u, H_\Omega u \rangle$ and evaluate it on functions u from the form core $C_0^\infty(\Omega)$. We get

$$\begin{aligned} \langle u, H_\Omega u \rangle_{L^2(\Omega)} &= \|\nabla u\|_{L^2(\Omega)}^2 - \int_\Omega V|u|^2 dx \\ &= \|\nabla' u\|_{L^2(\Omega)}^2 + \int_{\mathbb{R}^{d-1}} dx' \int_{\Omega(x')} \left(|\partial_t u(x', t)|^2 - V_{x'}(t) |u(x', t)|^2 \right) dt, \end{aligned}$$

where ∇' denotes the gradient in the first $(d-1)$ -coordinates.

For fixed $x' \in \mathbb{R}^{d-1}$ the functions $u(x', \cdot)$ belong to $C_0^\infty(\Omega(x'))$ and therefore to the form core of $W(x', V)$. It follows that

$$\langle u, H_\Omega u \rangle_{L^2(\Omega)} \geq \|\nabla' u\|_{L^2(\Omega)}^2 - \int_{\mathbb{R}^{d-1}} \langle u(x', \cdot), W(x', V)u(x', \cdot) \rangle_{L^2(\Omega(x'))} dx'. \quad (6)$$

To apply operator-valued Lieb-Thirring inequalities we need to extend these forms to \mathbb{R}^d . More precisely, we extend both sides of (6) by zero to $C_0^\infty(\mathbb{R}^d \setminus \partial\Omega)$ which is a form core of $(-\Delta_{\mathbb{R}^d \setminus \Omega}) \oplus H_\Omega$. This operator corresponds to the left-hand side of (6), while the semi-bounded form on the right-hand side is closed on the larger domain $H^1(\mathbb{R}^{d-1}, L^2(\mathbb{R}))$, where it corresponds to the operator

$$-\Delta' \otimes \mathbb{I} - W(x', V) \quad (7)$$

defined in $L^2(\mathbb{R}^{d-1}, L^2(\mathbb{R}))$. Due to the positivity of $(-\Delta_{\mathbb{R}^d \setminus \Omega})$ the variational principle implies

$$R_\sigma(V; \Omega) = \text{Tr} \left(-\Delta_{\mathbb{R}^d \setminus \Omega} \oplus H_\Omega \right)_-^\sigma \leq \text{Tr} \left(-\Delta' \otimes \mathbb{I} - W(x', V) \right)_-^\sigma. \quad (8)$$

Now we apply sharp Lieb-Thirring inequalities [LW00] to the Schrödinger operator (7) with operator-valued potential $W(x', V)$. For $\sigma \geq 3/2$ we obtain

$$\text{Tr} \left(-\Delta' \otimes \mathbb{I} - W(x', V) \right)_-^\sigma \leq L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \text{Tr} W(x', V)^{\sigma+(d-1)/2} dx' \quad (9)$$

and the claim follows from (8) and (9). \square

3. CONSTANT POTENTIALS

In this section we assume $V \equiv \Lambda > 0$ on quasi-bounded open sets $\Omega \subset \mathbb{R}^d$, $d \geq 2$. First we remark the following relations between the eigenvalue means. For $0 \leq \gamma < \sigma$ we have [AL78]

$$R_\sigma(\Lambda; \Omega) = \frac{1}{B(\gamma+1, \sigma-\gamma)} \int_0^\infty \tau^{\sigma-\gamma-1} R_\gamma((\Lambda-\tau)_+; \Omega) d\tau, \quad (10)$$

where B denotes the Beta-function. Hence one can use bounds or asymptotic results for R_γ to deduce the respective results for R_σ with $\sigma > \gamma \geq 0$. Conclusions from eigenvalue means of higher order to means of lower order are more cumbersome since eigenvalue means of lower order are less smooth. To derive uniform bounds on the counting function, that is, on $R_0(\Lambda; \Omega)$ one can make use of the estimate [Lap97]

$$R_0(\Lambda; \Omega) \leq (\tau\Lambda)^{-\sigma} R_\sigma((1+\tau)\Lambda; \Omega), \quad \tau > 0, \quad \Lambda > 0, \quad \sigma > 0, \quad (11)$$

and optimize the right hand side in $\tau > 0$. In general, sharp constants are lost but usually the correct order of growth in Λ is preserved.

In the following we consider specific domains with infinite volume. The discreteness of the spectrum of the Dirichlet Laplace operator defined on these domains can be deduced from the following sufficient condition [Ada70].

Lemma 3.1. *Let Ω be an open subset of \mathbb{R}^d and let Q be a cube with sides parallel to the coordinate axes. Let $\mu_{d-1}(Q, \Omega)$ denote the maximum of the $(d-1)$ -dimensional measure of $P(Q \setminus \Omega)$, where the maximum is taken over all projections P onto $(d-1)$ -dimensional faces of Q .*

Assume that for every $\epsilon > 0$ there exist $h \leq 1$ and $r \geq 0$ such that for every cube Q of side length h with sides parallel to the coordinate axes and with $Q \cap \{x \in \mathbb{R}^d : |x| > r\} \neq \emptyset$ we have

$$\frac{\mu_{d-1}(Q, \Omega)}{h^{d+1}} \geq \frac{1}{\epsilon}.$$

Then the embedding $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact.

In the following examples the trace of the operator $W(x', \Lambda)$ given in (5) can be calculated explicitly and we find that Proposition 2.1 yields sharp estimates on eigenvalue means.

3.1. Horn-shaped regions. First we consider horn-shaped regions, domains stretching to infinity along distinguished directions, see [vdB92a] for a general definition. These regions turn out to be of interest in different situations, see e.g. [Sim83, vdB84, DS92, vdB92a, MM06, Lun10]. They were introduced in [Sim83], where the semiclassical limit of the counting function was calculated for domains

$$\Omega_\nu = \{(x, y) \in \mathbb{R}^2 : |x| \cdot |y|^\nu \leq 1\}, \quad \nu \geq 1, \quad (12)$$

see Figure 1. Note that discreteness of the spectrum of H_{Ω_ν} can be deduced from Lemma 3.1.




FIGURE 1. The set Ω_2 .

In [GW10] it was shown that the methods introduced in Section 2 yield sharp upper bounds on the trace of the heat kernel of the Dirichlet Laplacian on various horn-shaped regions. Here we derive sharp bounds on eigenvalue means and order-sharp bounds on the counting function.

Let us recall the following asymptotic results from [Sim83]. For $\nu > 1$ the limit

$$R_0(\Lambda; \Omega_\nu) = \zeta(\nu) \left(\frac{2}{\pi}\right)^\nu \frac{\Gamma(\frac{\nu}{2} + 1)}{\sqrt{\pi} \Gamma(\frac{\nu+3}{2})} \Lambda^{(\nu+1)/2} + o\left(\Lambda^{(\nu+1)/2}\right), \quad \Lambda \rightarrow \infty,$$

holds, where $\zeta(\nu)$ denotes the Zeta function. Moreover, for $\nu = 1$

$$R_0(\Lambda; \Omega_1) = \frac{1}{\pi} \Lambda \ln \Lambda + o(\Lambda \ln \Lambda), \quad \Lambda \rightarrow \infty.$$

Applying (10) with $\gamma = 0$ we obtain for $\sigma > 0$ and $\nu > 1$

$$R_\sigma(\Lambda; \Omega_\nu) = \zeta(\nu) \left(\frac{2}{\pi}\right)^\nu \frac{B(\frac{\nu}{2} + 1, \sigma + 1)}{B(\sigma + \frac{\nu+3}{2}, \frac{1}{2})} \Lambda^{\sigma+(\nu+1)/2} + o\left(\Lambda^{\sigma+(\nu+1)/2}\right), \quad \Lambda \rightarrow \infty, \quad (13)$$

and for $\nu = 1$

$$R_\sigma(\Lambda; \Omega_1) = \frac{1}{\pi(\sigma + 1)} \Lambda^{\sigma+1} \ln \Lambda + o\left(\Lambda^{\sigma+1} \ln \Lambda\right), \quad \Lambda \rightarrow \infty. \quad (14)$$

In order to treat domains in higher dimensions we generalize the notions from [Sim83] and put

$$\Omega_\nu = \left\{ (x', x_d) \in \mathbb{R}^{d-1} \times \mathbb{R} : |x'| \cdot |x_d|^{\nu/(d-1)} \leq 1 \right\}, \quad d \geq 2, \quad \nu > 1.$$

For these domains of infinite volume an application of Proposition 2.1 yields sharp spectral estimates.

Theorem 3.2. *For $\sigma \geq 3/2$, $\nu > 1$, and all $\Lambda > 0$ the estimate*

$$R_\sigma(\Lambda; \Omega_\nu) \leq \frac{\zeta(\nu)}{2^{d-1}(d-1)} \left(\frac{2}{\pi}\right)^\nu \frac{\Gamma(\frac{\nu}{2} + 1)\Gamma(\sigma + 1)}{\Gamma(\frac{d+1}{2})\Gamma(\sigma + \frac{d+1+\nu}{2})} \Lambda^{\sigma+(d-1+\nu)/2}$$

holds.

Remark. For $d = 2$ we conclude that the bound

$$R_\sigma(\Lambda; \Omega_\nu) \leq \zeta(\nu) \left(\frac{2}{\pi}\right)^\nu \frac{B(\frac{\nu}{2} + 1, \sigma + 1)}{B(\sigma + \frac{\nu+3}{2}, \frac{1}{2})} \Lambda^{\sigma+(\nu+1)/2}$$

holds for $\sigma \geq 3/2$ and all $\Lambda > 0$. Comparing this bound with the asymptotic relation (13) we see that the estimate is sharp: For horn-shaped regions, just as well as for bounded domains, the leading term of the semiclassical limit yields a uniform upper bound.

Proof of Theorem 3.2. In this setting the section $\Omega_\nu(x')$ consists of one open interval

$$(-|x'|^{(1-d)/\nu}, |x'|^{(1-d)/\nu}).$$

Since $V \equiv \Lambda$, the trace of the operator-valued potential $W(x', \Lambda)$ defined in (5) can be evaluated explicitly. We find

$$\text{Tr}W(x', \Lambda) = \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{4|x'|^{2(1-d)/\nu}} \right)_+.$$

Applying Proposition 2.1 yields

$$\begin{aligned} R_\sigma(\Lambda; \Omega_\nu) &\leq L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{4|x'|^{2(1-d)/\nu}} \right)_+^{\sigma+(d-1)/2} dx' \Lambda^{\sigma+(d-1)/2} \\ &= L_{\sigma, d-1}^{cl} \omega_{d-1} \sum_{j \in \mathbb{N}} \int_0^\infty \left(1 - \frac{\pi^2 j^2}{4\Lambda r^{2(1-d)/\nu}} \right)_+^{\sigma+(d-1)/2} r^{d-2} dr \Lambda^{\sigma+(d-1)/2} \\ &= L_{\sigma, d-1}^{cl} \omega_{d-1} \zeta(\nu) \left(\frac{2}{\pi}\right)^\nu \frac{\nu B(\sigma + \frac{d+1}{2}, \frac{\nu}{2})}{2(d-1)} \Lambda^{\sigma+(d-1+\nu)/2}, \end{aligned}$$

where ω_{d-1} denotes the volume of the unit sphere in \mathbb{R}^{d-1} . We insert the identity

$$L_{\sigma, d-1}^{cl} \omega_{d-1} \frac{\nu B\left(\sigma + \frac{d+1}{2}, \frac{\nu}{2}\right)}{2(d-1)} = \frac{\Gamma(\sigma+1)\Gamma\left(\frac{\nu}{2}+1\right)}{2^{d-1}(d-1)\Gamma\left(\frac{d+1}{2}\right)\Gamma\left(\sigma + \frac{d+1+\nu}{2}\right)}$$

and arrive at the claimed estimate. \square

Now we apply (11) to deduce order-sharp bounds on the counting function.

Corollary 3.3. *For $\nu > 1$ and all $\Lambda > 0$ the estimate*

$$R_0(\Lambda; \Omega_\nu) \leq C_{d,\nu} \Lambda^{(d-1+\nu)/2}$$

holds with a constant

$$C_{d,\nu} \leq \frac{(d+\nu+2)^{(d+\nu+2)/2}}{3^{3/2}(d+\nu-1)^{(d+\nu-1)/2}} \frac{\zeta(\nu)}{2^{d-1}(d-1)} \left(\frac{2}{\pi}\right)^\nu \frac{\Gamma\left(\frac{\nu}{2}+1\right)\Gamma\left(\frac{5}{2}\right)}{\Gamma\left(\frac{d+1}{2}\right)\Gamma\left(\frac{d+\nu}{2}+2\right)}.$$

Proof. We use (11) with $\sigma = 3/2$ and apply Theorem 3.2 to obtain

$$\begin{aligned} R_0(\Lambda; \Omega_\nu) &\leq \frac{1}{(\tau\Lambda)^{3/2}} \frac{\zeta(\nu)}{2^{d-1}(d-1)} \left(\frac{2}{\pi}\right)^\nu \frac{\Gamma\left(\frac{\nu}{2}+1\right)\Gamma\left(\frac{5}{2}\right)}{\Gamma\left(\frac{d+1}{2}\right)\Gamma\left(\frac{d+\nu}{2}+2\right)} ((1+\tau)\Lambda)^{(d+\nu)/2+1} \\ &\leq \frac{(1+\tau)^{(d+\nu)/2+1}}{\tau^{3/2}} \frac{\zeta(\nu)}{2^{d-1}(d-1)} \left(\frac{2}{\pi}\right)^\nu \frac{\Gamma\left(\frac{\nu}{2}+1\right)\Gamma\left(\frac{5}{2}\right)}{\Gamma\left(\frac{d+1}{2}\right)\Gamma\left(\frac{d+\nu}{2}+2\right)} \Lambda^{(d-1+\nu)/2}. \end{aligned}$$

Minimizing in $\tau > 0$ yields $\tau_{\min} = 3/(d+\nu-1)$ and inserting this we obtain the claimed result. \square

Let us now consider the critical case $\nu = 1$ in dimension $d = 2$. Here the domain yields two equally strong singularities and we cannot distinguish one direction. However, choosing an intermediate direction we obtain a sharp estimate with a remainder term.

Theorem 3.4. *Let $\sigma \geq 3/2$. Then for $\Lambda \leq \pi^2/16$ we have $R_\sigma(\Lambda; \Omega_1) = 0$ and for $\Lambda > \pi^2/16$ the estimate*

$$R_\sigma(\Lambda; \Omega_1) \leq \frac{1}{\pi(\sigma+1)} \Lambda^{\sigma+1} \ln \Lambda + \frac{C}{\sigma+1} \Lambda^{\sigma+1}$$

holds with a constant

$$C < \frac{33 + 16 \ln\left(\frac{4}{\pi}\right)}{8\pi} < 1.47.$$

Remark. Again, comparing this inequality with the asymptotics (14), we see that the main term of the bound is sharp.

Proof of Theorem 3.4. Since the function $|\Omega_1(x)| = \frac{1}{x}$ has non-integrable singularities at zero and at infinity we have to choose a coordinate system $(x_1, x_2) \in \mathbb{R}^2$ rotated by $\frac{\pi}{4}$ with respect to the coordinate system $(x, y) \in \mathbb{R}^2$ which was used in definition (12). We get

$$\Omega_1(x_1) = \left\{ x_2 \in \mathbb{R} : |x_2| \leq \sqrt{|x_1|^2 + 2} \right\}$$

for $|x_1| \leq \sqrt{2}$ and

$$\Omega_1(x_1) = \left\{ x_2 \in \mathbb{R} : \sqrt{|x_1|^2 - 2} \leq |x_2| \leq \sqrt{|x_1|^2 + 2} \right\}$$

for $|x_1| > \sqrt{2}$. To simplify the following calculations and the resulting bound we confine ourselves to rough estimates which are nevertheless sufficient to prove the sharp constant in the leading term. First, we estimate $|\Omega_1(x_1)| \leq 4$ for $|x_1| \leq 2$ and

$$|\Omega_1(x_1)| \leq 2 \left(\sqrt{|x_1|^2 + 2} - \sqrt{|x_1|^2 - 2} \right) \leq \frac{4}{|x_1|} + \frac{2}{|x_1|^3}$$

for $|x_1| > 2$.

Suppose that $\Lambda \leq \pi^2/16$. Since $|\Omega_1(x_1)| \leq 4$ for all $x_1 \in \mathbb{R}$ we get

$$\text{Tr}W(x_1, \Lambda) = \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{|\Omega(x_1)|^2} \right)_+ = 0$$

for all $x_1 \in \mathbb{R}$. From Proposition 2.1 it follows that $R_\sigma(\Lambda; \Omega_1) = 0$ for $\Lambda \leq \pi^2/16$. On the other hand, if $\Lambda > \pi^2/16$ Proposition 2.1 implies

$$\begin{aligned} R_\sigma(\Lambda; \Omega_1) &\leq L_{\sigma,1}^{cl} \int_{\mathbb{R}} \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{|\Omega(x_1)|^2} \right)_+^{\sigma+1/2} dx_1 \\ &\leq 4L_{\sigma,1}^{cl} \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{16} \right)_+^{\sigma+1/2} + 2L_{\sigma,1}^{cl} \int_2^\infty \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{l(x_1)^2} \right)_+^{\sigma+1/2} dx_1, \end{aligned} \quad (15)$$

with

$$l(x_1) = \frac{4}{|x_1|} + \frac{2}{|x_1|^3}.$$

Note that for $A > 0$ and $\gamma > 0$ we have

$$\sum_{j \in \mathbb{N}} \left(1 - \frac{j^2}{A^2} \right)_+^\gamma \leq \frac{A}{2} B\left(\frac{1}{2}, \gamma + 1\right), \quad (16)$$

thus

$$4L_{\sigma,1}^{cl} \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{16} \right)_+^{\sigma+1/2} \leq \frac{8}{\pi} L_{\sigma,1}^{cl} B\left(\frac{1}{2}, \sigma + \frac{3}{2}\right) \Lambda^{\sigma+1} = \frac{4}{\pi} \frac{1}{\sigma + 1} \Lambda^{\sigma+1}. \quad (17)$$

Now we turn to the second term in (15). Put $x(\Lambda) = (4\sqrt{\Lambda})/\pi + \pi/(4\sqrt{\Lambda})$. For $x_1 \geq x(\Lambda)$ we have $l(x_1) \leq \pi/\sqrt{\Lambda}$, hence

$$\sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{l(x_1)^2} \right)_+ = 0.$$

In view of (16) it follows that

$$\begin{aligned} 2L_{\sigma,1}^{cl} \int_2^\infty \sum_{j \in \mathbb{N}} \left(\Lambda - \frac{\pi^2 j^2}{l(x_1)^2} \right)_+^{\sigma+1/2} dx_1 &\leq \frac{1}{\pi} L_{\sigma,1}^{cl} B\left(\frac{1}{2}, \sigma + \frac{3}{2}\right) \int_2^{x(\Lambda)} l(x_1) dx_1 \Lambda^{\sigma+1} \\ &= \frac{1}{2\pi} \frac{1}{\sigma + 1} \int_2^{x(\Lambda)} l(x_1) dx_1 \Lambda^{\sigma+1}. \end{aligned} \quad (18)$$

By definition of $x(\Lambda)$ and $l(x_1)$,

$$\begin{aligned} \int_2^{x(\Lambda)} l(x_1) dx_1 &= \int_2^{x(\Lambda)} \left(\frac{4}{x_1} + \frac{2}{x_1^3} \right) dx_1 \\ &\leq 2 \ln \Lambda + 4 \ln \left(\frac{4}{\pi} + \frac{\pi}{4\Lambda} \right) - 4 \ln 2 + \frac{1}{4} \\ &\leq 2 \ln \Lambda + 4 \ln \left(\frac{4}{\pi} \right) + \frac{1}{4} \end{aligned} \tag{19}$$

for $\Lambda > \pi^2/16$. Inserting (17), (18) and (19) into (15) finishes the proof. \square

Again we can apply (11) to deduce order-sharp bounds on the counting function.

Corollary 3.5. *For $\Lambda \leq \pi^2/16$ we have $R_0(\Lambda; \Omega_1) = 0$ and for $\Lambda > \pi^2/16$ the estimate*

$$R_0(\Lambda; \Omega_1) \leq \left(\frac{5}{3} \right)^{3/2} \frac{1}{\pi} \Lambda \ln \Lambda + C \Lambda,$$

holds, with a constant

$$C < \sqrt{\frac{5}{3}} \frac{825 + 400 \ln \left(\frac{4}{\pi} \right) + 360\pi \ln \left(\frac{5}{3} \right)}{72\pi} < 8.56.$$

3.2. Spiny urchins. In this subsection we study the eigenvalues of the Dirichlet Laplacian on so called spiny urchins, radially symmetric domains $\Omega_S \subset \mathbb{R}^2$ with infinite volume, which were introduced in [Cla67].

To construct Ω_S we use polar coordinates $(r, \varphi) \in [0, \infty) \times [0, 2\pi)$ and choose an increasing sequence $(r_n)_{n \in \mathbb{N}}$ of positive real numbers and put $r_0 = 0$. For $n \in \mathbb{N}_0$ and $k = 1, 2, \dots, 2^{n+2}$ let

$$\Gamma_{n,k} = \left\{ (r, \varphi) : r \geq r_n, \varphi = \frac{k-1}{2^{n+1}} \pi \right\}$$

be semi-axes and define

$$\Omega_S = \mathbb{R}^2 \setminus \bigcup_{n,k} \Gamma_{n,k},$$

see Figure 2. Note that this domain, though quasi bounded, has empty exterior. However, if

$$\lim_{n \rightarrow \infty} r_n 2^{-n} = 0, \tag{20}$$

then discreteness of the spectrum of H_{Ω_S} can be deduced from Lemma 3.1, see also [vdB92b].

For $r_n = n$ the domain Ω_S was analyzed in [Fle78], where the leading term of the semiclassical limit was calculated: For $r_n = n$ the asymptotic relation


$$R_0(\Lambda; \Omega_S) = C \Lambda (\ln \Lambda)^2 + o(\Lambda (\ln \Lambda)^2), \quad \Lambda \rightarrow \infty,$$

holds with a constant $C > 0$.

The general setting of an arbitrary increasing sequence $(r_n)_{n \in \mathbb{N}_0}$ was studied in [vdB92b]: If $r_0 > 0$ and (20) is satisfied then for all $\Lambda > 2^{14} r_0^{-2}$ the bound

$$R_0(\Lambda; \Omega_S) \leq 50(8^{-1} + 8\pi)^2 \Lambda r_{K(\Lambda)}^2$$

holds with $K(\Lambda) = \max\{n \in \mathbb{N} : r_n 2^{-n} > (32)^{-1} \sqrt{\Lambda}\}$. Moreover, there is a similar lower bound. Here we extend and improve the upper bound: We derive order-sharp estimates on the eigenvalue means of H_{Ω_S} valid for all $\Lambda > 0$.

FIGURE 2. The set Ω_S .

First, we need to adapt Proposition 2.1 to the radially symmetric situation. For $r \in (0, \infty)$ put

$$\Omega_S(r) = \{\varphi \in [0, 2\pi) : (r, \varphi) \in \Omega_S\} .$$

Then $\Omega_S(r)$ consists of finitely many open intervals $I_k(r)$, $k = 1, \dots, N(r)$. Choose $u \in C_0^\infty(\Omega_S)$ and consider the quadratic form

$$\begin{aligned} \langle u, H_{\Omega_S} u \rangle_{L^2(\Omega_S)} &= \int_{\Omega} \overline{u(x)} (-\Delta u(x) - \Lambda u(x)) dx \\ &= \int_0^\infty \int_{\Omega_S(r)} \overline{u(r, \varphi)} \left(-\partial_r^2 - \frac{1}{r} \partial_r - \frac{1}{r^2} \partial_\varphi^2 - \Lambda \right) u(r, \varphi) d\varphi r dr . \end{aligned} \quad (21)$$

For fixed $r > 0$ the function $u_r(\varphi) = u(r, \varphi)$ belongs to $C_0^\infty(\Omega_S(r))$. It satisfies Dirichlet boundary conditions at the endpoints of the intervals $I_k(r)$, $k = 1, \dots, N(r)$.

To rewrite the form in the ground state representation put $v(r, \varphi) = \sqrt{r} u(r, \varphi)$. Then again $v(r, \varphi)$ belongs to $C_0^\infty(\Omega_S)$ and for fixed $r > 0$, we have $v_r(\varphi) = v(r, \varphi) \in C_0^\infty(\Omega_S(r))$. Moreover,

$$\int_0^\infty \int_{\Omega_S(r)} |u(r, \varphi)|^2 d\varphi r dr = \int_0^\infty \int_{\Omega_S(r)} |v(r, \varphi)|^2 d\varphi dr$$

and

$$\left(-\partial_r^2 - \frac{1}{r} \partial_r - \frac{1}{r^2} \partial_\varphi^2 \right) u(r, \varphi) = \frac{1}{\sqrt{r}} \left(-\partial_r^2 - \frac{1}{4r^2} - \frac{1}{r^2} \partial_\varphi^2 \right) v(r, \varphi) .$$

Inserting this into (21) we obtain

$$\langle u, H_{\Omega_S} u \rangle_{L^2(\Omega_S)} = \int_0^\infty \int_{\Omega_S(r)} \left(|\partial_r v|^2 + \frac{1}{r^2} |\partial_\varphi v|^2 - \left(\frac{1}{4r^2} + \Lambda \right) |v|^2 \right) d\varphi dr . \quad (22)$$

In this setting, we define the Schrödinger-type operators

$$H_k(r) = -\frac{1}{r^2} \frac{d^2}{d\varphi^2} - \left(\frac{1}{4r^2} + \Lambda \right), \quad k = 1, \dots, N(r),$$

in $L^2(I_k(r))$ with Dirichlet boundary conditions at the endpoints of $I_k(r)$. In the same way as in (5) let

$$W(r, \Lambda) = \bigoplus_{k=1}^{N(r)} H_k(r)_-$$

be the negative part of the operator

$$-\frac{1}{r^2} \frac{d^2}{d\varphi^2} - \left(\frac{1}{4r^2} + \Lambda \right)$$

in $L^2(\Omega_S(r))$ with Dirichlet boundary conditions. In view of (22) we can apply Proposition 2.1 and for $\sigma \geq 3/2$ we get

$$R_\sigma(\Lambda; \Omega_S) \leq L_{\sigma,1}^{cl} \int_0^\infty \text{Tr} W(r, \Lambda)^{\sigma+1/2} dr. \quad (23)$$

To estimate the right hand side and to derive bounds on the eigenvalues means we assume that (20) is satisfied and that

$$r_{n+1} \leq 2r_n \quad (24)$$

holds for all $n \in \mathbb{N}$. Then the sequence

$$\frac{2^{2n}}{r_n^2} - \frac{1}{4r_n^2}, \quad n \in \mathbb{N},$$

is increasing and for all $\Lambda > 15/4 \cdot r_1^{-2}$ there is a unique index $\hat{n}(\Lambda) \in \mathbb{N}$ satisfying

$$\Lambda > \frac{2^{2n}}{r_n^2} - \frac{1}{4r_n^2} \text{ for all } n \leq \hat{n}(\Lambda) \quad \text{and} \quad \Lambda \leq \frac{2^{2n}}{r_n^2} - \frac{1}{4r_n^2} \text{ for all } n > \hat{n}(\Lambda). \quad (25)$$

To simplify notation we put $\hat{r}(\Lambda) = r_{\hat{n}(\Lambda)}$.

Lemma 3.6. *Let $\sigma \geq 3/2$ and assume that (20) and (24) are satisfied. Then for $\Lambda \leq 15/4 \cdot r_1^{-2}$ we have $R_\sigma(\Lambda; \Omega_S) = 0$ and for $\Lambda > 15/4 \cdot r_1^{-2}$ the estimate*

$$R_\sigma(\Lambda; \Omega_S) \leq L_{\sigma,2}^{cl} \pi \hat{r}(\Lambda)^2 \Lambda^{\sigma+1} + C_\sigma \Lambda^\sigma \ln(\Lambda \hat{r}(\Lambda)).$$

holds with a constant $C_\sigma > 0$ depending only on σ .

Remark. If we compare the main term of this bound with the Berezin inequality (4) we see that the effective domain that enters into the bound is a disk with radius $\hat{r}(\Lambda)$.

Proof of Lemma 3.6. In view of (23) we have to estimate

$$\text{Tr} W(r, \Lambda) = \text{Tr} \left(-\frac{1}{r^2} \frac{d^2}{d\varphi^2} - \Lambda - \frac{1}{4r^2} \right)_- = \sum_{k=1}^{N(r)} \sum_{j \in \mathbb{N}} \left(\Lambda + \frac{1}{4r^2} - \frac{\pi^2 j^2}{r^2 |I_k(r)|^2} \right)_+.$$

Fix $r > 0$ and $n_0 \in \mathbb{N}$ such that $r_{n_0-1} < r \leq r_{n_0}$. Then the section $\Omega_S(r) \subset [0, 2\pi)$ consists of 2^{n_0+1} identical open intervals of length $|I_k(r)| = \pi/2^{n_0}$. Hence,

$$\text{Tr} W(r, \Lambda) = 2^{n_0+1} \sum_{j \in \mathbb{N}} \left(\Lambda + \frac{1}{4r^2} - \frac{2^{2n_0} j^2}{r^2} \right)_+.$$

Note that for all $j \in \mathbb{N}$

$$\frac{2^{2n_0} j^2}{r^2} - \frac{1}{4r^2} \geq \frac{2^{2n_0+2} - 1}{4r^2} \geq \frac{15}{4r_1^2}.$$

For $\Lambda \leq 15/4 \cdot r_1^{-2}$ we obtain $\text{Tr}W(r, \Lambda) = 0$ and by (23) also $R_\sigma(\Lambda; \Omega_S) = 0$.

Hence, we can assume $\Lambda > 15/4 \cdot r_1^{-2}$. Suppose that $r > \hat{r}(\Lambda)$ thus $n_0 > \hat{n}(\Lambda)$. From (25) we get

$$\frac{2^{2n_0} j^2}{r^2} - \frac{1}{4r^2} \geq \frac{2^{2n_0+2} - 1}{4r_{n_0}^2} \geq \Lambda$$

for all $j \in \mathbb{N}$ and it follows that $\text{Tr}W(r, \Lambda) = 0$ for $r > \hat{r}(\Lambda)$. Moreover, if $r^2 \leq 15/(4\Lambda)$ we have $r \leq r_1$ and

$$\frac{4j^2}{r^2} - \frac{1}{4r^2} \geq \frac{15}{4r^2} \geq \Lambda$$

for all $j \in \mathbb{N}$. Again it follows that $\text{Tr}W(r, \Lambda) = 0$ and it remains to consider $\sqrt{15}/(2\sqrt{\Lambda}) < r < \hat{r}(\Lambda)$.

For such r we apply (16) to estimate

$$\text{Tr}W(r, \Lambda)^{\sigma+1/2} = 2^{n_0+1} \sum_{j \in \mathbb{N}} \left(\Lambda + \frac{1}{4r^2} - \frac{2^{2n_0} j^2}{r^2} \right)_+^{\sigma+1/2} \leq r \left(\Lambda + \frac{1}{4r^2} \right)^{\sigma+1} B\left(\frac{1}{2}, \sigma + \frac{3}{2}\right).$$

From (23) we conclude

$$\begin{aligned} R_\sigma(\Lambda; \Omega_S) &\leq L_{\sigma,1}^{\text{cl}} B\left(\frac{1}{2}, \sigma + \frac{3}{2}\right) \int_{\sqrt{15}/(2\sqrt{\Lambda})}^{\hat{r}(\Lambda)} r \left(\Lambda + \frac{1}{4r^2} \right)^{\sigma+1} dr \\ &= \frac{1}{16(\sigma+1)} \Lambda^\sigma \int_{15}^{4\Lambda\hat{r}(\Lambda)^2} \left(1 + \frac{1}{s} \right)^{\sigma+1} ds \\ &\leq \frac{1}{4(\sigma+1)} \hat{r}(\Lambda)^2 \Lambda^{\sigma+1} + \frac{16^{\sigma-1}}{15^\sigma} \Lambda^\sigma \ln(4\Lambda\hat{r}(\Lambda)^2) \end{aligned}$$

and the claim of the lemma follows from the identity $4\pi(\sigma+1)L_{\sigma,2}^{\text{cl}} = 1$. \square

Before we give examples we supplement Lemma 3.6 with the following lower bound on $R_\sigma(\Lambda; \Omega_S)$.

Lemma 3.7. *Assume there exists $N_0 \in \mathbb{N}$ such that $r_{n-1} < (1 - 2^{-n})r_n$ is satisfied for all $n \geq N_0$. Then for $\sigma \geq 0$ there exist positive constants C and μ independent of Λ such that*

$$R_\sigma(\Lambda; \Omega_S) \geq C \sum_{n=N_0}^{\hat{n}(\mu\Lambda)} r_n (r_n - r_{n-1}) \Lambda^{\sigma+1}$$

holds for $\Lambda > 0$ with $\hat{n}(\mu\Lambda) > N_0$.

Proof. For $n \geq N_0$ and $k \in \{1, \dots, 2^{n+1}\}$ consider a segment $\Omega_{n,k} \subset \Omega_S$, i.e., a region between $r = r_{n-1}$, $r = r_n$ and two adjacent semi-axes $\Gamma_{n,k}$ and $\Gamma_{n,k+1}$. Note that there are 2^{n+1} identical segments $\Omega_{n,k}$. Let $\tau(n)$ denote the maximal number of disjoint squares Q_{l_n}

with side length $l_n = r_n/2^{n+1}$ that can be placed in the interior of $\Omega_{n,k}$. From the definition of Ω_S it follows that

$$\tau(n) \geq C \frac{r_n - r_{n-1}}{l_n}, \quad n \geq N_0.^1$$

Hence, the variational principle implies

$$R_\sigma(\Lambda; \Omega_S) \geq \sum_{n \geq N_0} 2^{n+1} \tau(n) R_\sigma(\Lambda; Q_{l_n}) \geq C \sum_{n \geq N_0} 2^{n+1} \frac{r_n - r_{n-1}}{l_n} R_\sigma(\Lambda; Q_{l_n}). \quad (26)$$

To estimate $R(\Lambda; Q_{l_n})$ from below, we first consider the square Q_1 with side length 1. From Weyl's asymptotic law (3) we know that there are positive constants C and Λ_0 , such that $R_\sigma(\Lambda; Q_1) \geq C \Lambda^{\sigma+1}$ holds for all $\Lambda \geq \Lambda_0$. By scaling, we deduce that

$$R_\sigma(\Lambda; Q_{l_n}) \geq C l_n^2 \Lambda^{\sigma+1} \quad (27)$$

holds for all $\Lambda \geq \Lambda_0/l_n^2$.

Fix $\Lambda > 0$. From (25) we deduce that

$$\frac{\Lambda_0}{l_n^2} = 4\Lambda_0 \frac{2^{2n}}{r_n^2} \leq 8\Lambda_0 \left(\frac{2^{2n}}{r_n^2} - \frac{1}{4r_n^2} \right) \leq \Lambda$$

holds if $n \leq \hat{n}(\Lambda/(8\Lambda_0))$. Denoting $\mu = 1/(8\Lambda_0)$ we find that (27) is valid for all squares Q_{l_n} with $n \leq \hat{n}(\mu\Lambda)$.

In view of (26) it follows that

$$R_\sigma(\Lambda; \Omega_S) \geq C \sum_{n=N_0}^{\hat{n}(\mu\Lambda)} 2^{n+1} \frac{r_{n+1} - r_n}{l_n} l_n^2 \Lambda^{\sigma+1} \geq C \sum_{n=N_0}^{\hat{n}(\mu\Lambda)} r_n (r_n - r_{n-1}) \Lambda^{\sigma+1}$$

and the proof is complete. \square

Let us state some examples to show that the bounds capture the correct order in Λ and that choosing different sequences $(r_n)_{n \in \mathbb{N}}$ leads to different behavior in the semiclassical limit.

Corollary 3.8. *Let $\sigma \geq 0$.*

(1) *Assume $r_n = n$. Then for $0 < \Lambda \leq 15/4$ we have $R_\sigma(\Lambda; \Omega_S) = 0$ and for $\Lambda > 15/4$*

$$R_\sigma(\Lambda; \Omega_S) \leq C_\sigma \Lambda^{\sigma+1} (\ln \Lambda)^2.$$

(2) *Assume $r_n = 2^{\delta n}$ with $0 < \delta < 1$. Then for $0 < \Lambda \leq 15 \cdot 2^{-2(1+\delta)}$ we have $R_\sigma(\Lambda; \Omega_S) = 0$ and for $\Lambda > 15 \cdot 2^{-2(1+\delta)}$*

$$R_\sigma(\Lambda; \Omega_S) \leq C_{\sigma,\delta} \Lambda^{\sigma+1/(1-\delta)}.$$

All bounds capture the correct order in Λ as $\Lambda \rightarrow \infty$.

Proof. To prove the bounds for $\sigma \geq 3/2$ we can apply Lemma 3.6 and it remains to estimate $\hat{r}(\Lambda)$. By definition, $\hat{r}(\Lambda) = r_{\hat{n}(\Lambda)}$ and by (25) $r_{\hat{n}(\Lambda)}$ satisfies

$$\frac{2^{2\hat{n}(\Lambda)}}{r_{\hat{n}(\Lambda)}^2} - \frac{1}{4r_{\hat{n}(\Lambda)}^2} \leq \Lambda.$$

It follows that $\hat{r}(\Lambda) \leq C \ln \Lambda$ in the case $r_n = n$ and $\hat{r}(\Lambda) \leq C_\delta \Lambda^{\delta/(2(1-\delta))}$ in the case $r_n = 2^{\delta n}$ and the bounds for $\sigma \geq 3/2$ follow from Lemma 3.6. To deduce the claimed estimates for $0 \leq \sigma < 3/2$ we apply (11) and finally (10).

¹Here and in the following the letter C denotes various positive constants that are independent of Λ .

It remains to prove that the estimates are of correct order in Λ . Note that in the case $r_n = n$ the assumptions of Lemma 3.7 are satisfied with $N_0 = 1$. Hence, we have

$$\sum_{n=N_0}^{\hat{n}(\mu\Lambda)} r_n (r_n - r_{n-1}) = \sum_{n=1}^{\hat{n}(\mu\Lambda)} n \geq C\hat{n}(\mu\Lambda)^2 = C\hat{r}(\mu\Lambda)^2.$$

In the case $r_n = 2^{\delta n}$ we find for sufficiently large Λ that

$$\sum_{n=N_0}^{\hat{n}(\mu\Lambda)} r_n (r_n - r_{n-1}) = \sum_{n=N_0}^{\hat{n}(\mu\Lambda)} 2^{\delta n} (2^{\delta n} - 2^{\delta(n-1)}) \geq C \sum_{n=N_0}^{\hat{n}(\mu\Lambda)} 2^{2\delta n} \geq C2^{2\delta\hat{n}(\mu\Lambda)} = C\hat{r}(\mu\Lambda)^2,$$

holds. In both cases, we insert this into Lemma 3.7 and get

$$R_\sigma(\Lambda; \Omega_S) \geq C\Lambda^{\sigma+1} \hat{r}(\mu\Lambda)^2. \quad (28)$$

For Λ large enough the relations (25) imply

$$\hat{r}(\mu\Lambda) \geq C \ln(\mu\Lambda) \geq C \ln \Lambda$$

if $r_n = n$ and

$$\hat{r}(\mu\Lambda) \geq C(\mu\Lambda)^{\delta/(2(1-\delta))} \geq C\Lambda^{\delta/(2(1-\delta))}$$

if $r_n = 2^{\delta n}$. As $\Lambda \rightarrow \infty$ we obtain from (28) that $R_\sigma(\Lambda; \Omega_S) = O(\Lambda^{\sigma+1}(\ln \Lambda)^2)$ in the case $r_n = n$ and $R_\sigma(\Lambda; \Omega_S) = O(\Lambda^{\sigma+1/(1-\delta)})$ in the case $r_n = 2^{\delta n}$. Thus the bounds on $R_\sigma(\Lambda, \Omega_S)$ show the correct order in Λ . \square

Let us state one more example, where one encounters exponential growth of the eigenvalue means.

Corollary 3.9. *Assume $\sigma \geq 3/2$ and $r_n = 2^n/\sqrt{n}$. Then for $0 < \Lambda < 15/16$ we have $R_\sigma(\Lambda; \Omega_S) = 0$ and for $\Lambda > 15/16$*

$$R_\sigma(\Lambda; \Omega_S) \leq C_\sigma 2^{2\Lambda} \Lambda^\sigma.$$

This bound follows from Lemma 3.6 similar as in Corollary 3.8.

4. NON-CONSTANT POTENTIALS

In this section we consider Schrödinger operators H_Ω with non-constant potentials $V \geq 0$ on open sets $\Omega \subset \mathbb{R}^d$. Since we define H_Ω with Dirichlet boundary conditions the variational principle implies that the sharp Lieb-Thirring inequality (2) holds. In fact, the Dirichlet condition gives rise to an improvement of this bound. In this section we use this to derive sharp Lieb-Thirring inequalities with remainder term.

4.1. One-dimensional considerations. As in Section 3 we can apply Proposition 2.1 to reduce the problem to one dimension. However, for non-constant potentials V the trace of the operator-valued potential $W(x', V)$ defined in (5) cannot be calculated explicitly. Therefore we need to study the one-dimensional situation in more detail to derive the following improvement of (2).

Theorem 4.1. *Let $I \subset \mathbb{R}$ be an open interval of length $l < \infty$ and assume $\sigma \geq 3/2$ and $V \in L^{\sigma+1/2}(I)$ such that*

$$A = l \int_I V(t) dt < \infty.$$

Then for $A \leq 2 \ln 3$ we have $R_\sigma(V; I) = 0$ and for $A > 2 \ln 3$

$$R_\sigma(V; I) \leq L_{\sigma,1}^{cl} \int_I V(t)^{\sigma+1/2} dt - \left(\frac{2 \left(\int_I V(t) dt \right)^2}{\exp(A) - 1} \right)^\sigma.$$

The remainder of Section 4.1 is devoted to the proof of this result. In particular, we study the effect of different boundary conditions on the eigenvalues. First we assume $I = (0, l)$ and $V \in C_0^\infty(I)$. Recall that

$$H_I = -\frac{d^2}{dt^2} - V$$

is defined in $L^2(I)$ as self-adjoint operator generated by the quadratic form

$$\langle u, H_I u \rangle = \int (|u'(t)|^2 - V(t)|u(t)|^2) dt, \quad (29)$$

with form domain $H_0^1(I)$. Moreover, we define the operator

$$H_{\mathbb{R}} = -\frac{d^2}{dt^2} - V$$

in $L^2(\mathbb{R})$ generated by the form (29) with form domain $H^1(\mathbb{R})$.

We assume that the negative spectrum of H_I consists of N eigenvalues $(-\lambda_k)_{k=1}^N$, $N \in \mathbb{N}$, and denote the negative eigenvalues of $H_{\mathbb{R}}$ by $(-\mu_k)_{k=1}^M$. The variational principle implies $M \geq N$ and $-\mu_k \leq -\lambda_k$ for each $k = 1, \dots, N$.

In order to derive relations between the eigenvalues of H_I and $H_{\mathbb{R}}$ we define

$$H_I^{(\alpha, \beta)} = -\frac{d^2}{dt^2} - V, \quad 0 \leq \alpha, \beta \leq \frac{\pi}{2},$$

as self-adjoint operators generated by the form

$$\langle u, H_I^{(\alpha, \beta)} u \rangle = \int |u'(t)|^2 dt - \int V(t)|u(t)|^2 dt + (\cot \alpha) |u(0)|^2 + (\cot \beta) |u(l)|^2$$

with form domain $H^1(I)$. Note that eigenfunctions of $H_I^{(\alpha, \beta)}$ satisfy boundary conditions of the third kind: $u'(0) = (\cot \alpha)u(0)$ and $u'(l) = -(\cot \beta)u(l)$. For $\alpha, \beta \in [0, \frac{\pi}{2}]$ the negative spectrum of $H_I^{(\alpha, \beta)}$ consists of eigenvalues $(-\nu_k(\alpha, \beta))_{k=1}^{N(\alpha, \beta)}$. We point out that for $\alpha = \beta = 0$ we recover Dirichlet boundary conditions:

$$H_I^{(0,0)} = H_I, \quad N(0,0) = N, \quad \text{and} \quad (\nu_k(0,0))_{k=1}^{N(0,0)} = (\lambda_k)_{k=1}^N. \quad (30)$$

We need the following result from [Wei03] about the behavior of the eigenvalues of $H_I^{(\alpha, \beta)}$. For $\alpha \in [0, \frac{\pi}{2}]$ and $\nu > 0$ let $u(t; \nu, \alpha)$ to be the unique solution of

$$\begin{aligned} -u''(t) - V(t)u(t) &= -\nu u(t), \quad t \in I, \\ u(0; \nu, \alpha) &= \sin \alpha, \\ u'(0; \nu, \alpha) &= \cos \alpha. \end{aligned} \quad (31)$$

Lemma 4.2. Fix $\beta \in [0, \frac{\pi}{2}]$. Then for $\alpha \in (0, \frac{\pi}{2})$ the map $\alpha \mapsto \nu_k(\alpha, \beta)$ is monotone increasing and differentiable and we have

$$\frac{d\nu_k(\alpha, \beta)}{d\alpha} = \|u(\cdot; \nu_k(\alpha, \beta), \alpha)\|_{L^2(I)}^{-2}.$$

Because of the symmetry of the eigenvalue problem (31) a corresponding result holds for fixed $\alpha \in [0, \frac{\pi}{2}]$ and the map $\beta \mapsto \nu_k(\alpha, \beta)$, $\beta \in [0, \frac{\pi}{2}]$. For $k = 1, \dots, N$ it follows that

$$-\nu_k(\alpha, \alpha) \leq -\nu_k(0, 0) = -\lambda_k < 0$$

for all $\alpha \in [0, \frac{\pi}{2}]$.

For $k = 1, \dots, N$ put

$$\omega_k = \operatorname{arccot} \sqrt{\mu_k} \in \left[0, \frac{\pi}{2}\right]. \quad (32)$$

Then we have $N(\omega_k, \omega_k) \geq N$ and both $-\mu_k$ and $-\nu_k(\omega_k, \omega_k)$ exist as negative eigenvalues of $H_{\mathbb{R}}$ and $H_I^{(\omega_k, \omega_k)}$ respectively.

Proposition 4.3. *For $k = 1, \dots, N$ the eigenvalues of $H_{\mathbb{R}}$ and $H_I^{(\omega_k, \omega_k)}$ satisfy*

$$-\mu_k = -\nu_k(\omega_k, \omega_k).$$

Proof. For arbitrary $k \in \{1, \dots, N\}$ let Φ_k denote the eigenfunction of $H_{\mathbb{R}}$ corresponding to $-\mu_k$. Then $\operatorname{supp} V \subset I = (0, l)$ implies

$$\begin{aligned} \Phi_k(t) &= c_1 \exp(-\sqrt{\mu_k}t) \quad \text{for } t \geq l \quad \text{and} \\ \Phi_k(t) &= c_2 \exp(+\sqrt{\mu_k}t) \quad \text{for } t \leq 0 \end{aligned}$$

with suitable constants $c_1, c_2 \in \mathbb{R}$. From (32) it follows that $\Phi_k'(0) = (\cot \omega_k)\Phi_k(0)$ and $\Phi_k'(l) = -(\cot \omega_k)\Phi_k(l)$. Put $\tilde{\Phi}_k = \Phi_k|_{(0, l)}$. Since $\tilde{\Phi}_k$ belongs to the domain of $H_I^{(\omega_k, \omega_k)}$ we find that $-\mu_k$ is an eigenvalue of $H_I^{(\omega_k, \omega_k)}$. Note that Φ_k has $k - 1$ zeros in the interior of I . Therefore $\tilde{\Phi}_k$ has $k - 1$ zeros as well and we conclude $-\mu_k = -\nu_k(\omega_k, \omega_k)$. \square

Similar as in (31) let $\tilde{u}(t; \nu, \beta)$, $\beta \in [0, \frac{\pi}{2}]$, $\nu > 0$, be the unique solution of

$$\begin{aligned} -\tilde{u}''(t) - V(t)\tilde{u}(t) &= -\nu \tilde{u}(t), \quad t \in I, \\ \tilde{u}(l; \nu, \beta) &= \sin \beta, \\ \tilde{u}'(l; \nu, \beta) &= -\cos \beta. \end{aligned}$$

Due to the symmetry of the eigenvalue problem (31) there is a result analogous to Lemma 4.2 relating the derivative of the map $\beta \mapsto \nu_k(\alpha, \beta)$ to the L^2 -norm of $\tilde{u}(\cdot; \nu_k(\alpha, \beta), \beta)$.

In view of (30) and Proposition 4.3 we have

$$\mu_k - \lambda_k = \nu_k(\omega_k, \omega_k) - \nu_k(0, 0) = \nu_k(\omega_k, \omega_k) - \nu_k(0, \omega_k) + \nu_k(0, \omega_k) - \nu_k(0, 0).$$

Hence, applying Lemma 4.2 and its analog for the map $\beta \mapsto \nu_k(0, \beta)$ yields

$$\mu_k - \lambda_k = \int_0^{\omega_k} \|u(\cdot; \nu_k(\alpha, \omega_k), \alpha)\|_{L^2(I)}^{-2} d\alpha + \int_0^{\omega_k} \|\tilde{u}(\cdot; \nu_k(0, \beta), \beta)\|_{L^2(I)}^{-2} d\beta \quad (33)$$

for $k = 1, \dots, N$.

In the remainder of this subsection we use this identity to complete the proof of Theorem 4.1. In order to get a result valid without further assumptions on the potential V we have to restrict ourselves to considering the ground states.

Lemma 4.4. *Let $I \subset \mathbb{R}$ be an open interval of length l and $V \in C_0^\infty(I)$. Then the inequality*

$$\mu_1 - \lambda_1 \geq \frac{2 \left(\int V(t) dt \right)^2}{\exp \left(l \int V(t) dt \right) - 1}$$

holds. Moreover, if $l \int V(t) dt \leq 2 \ln 3$ then $-\lambda_1 \geq 0$ and we have $R_\sigma(V; I) = 0$ for $\sigma \geq 0$.

Proof. First we remark that it suffices to prove the result for $I = (0, l)$. To apply (33) we have to analyze the functions $u(\cdot; \nu_1(\alpha, \omega_1), \alpha)$ and $\tilde{u}(\cdot; \nu_1(0, \beta), \beta)$ for $0 < \alpha, \beta < \omega_1$.

By definition, the function u is the first eigenfunction of $H_I^{(\alpha, \omega_1)}$ thus it is non-negative on I . As a solution of (31) u solves the integral equation

$$\begin{aligned} u(t; \nu, \alpha) &= \frac{1}{2} (\sin \alpha) \left(e^{\sqrt{\nu}t} + e^{-\sqrt{\nu}t} \right) + \frac{1}{2} (\cos \alpha) \left(\frac{e^{\sqrt{\nu}t} - e^{-\sqrt{\nu}t}}{\sqrt{\nu}} \right) \\ &\quad - \int_0^t \frac{\sinh(\sqrt{\nu}(t-s))}{\sqrt{\nu}} V(s) u(s; \nu, \alpha) ds. \end{aligned} \quad (34)$$

The first two summands are non-decreasing in $\nu > 0$. For $\alpha \in [0, \omega_1]$, Lemma 4.2 and Proposition 4.3 imply $\nu_1(\alpha, \omega_1) \leq \mu_1$. Since the integrand in (34) is positive it follows that

$$\begin{aligned} u(t; \nu_1(\alpha, \omega_1), \alpha) &\leq \frac{1}{2} (\sin \alpha) \left(e^{\sqrt{\mu_1}t} + e^{-\sqrt{\mu_1}t} \right) + \frac{1}{2} (\cos \alpha) \left(\frac{e^{\sqrt{\mu_1}t} - e^{-\sqrt{\mu_1}t}}{\sqrt{\mu_1}} \right) \\ &= \frac{1}{2} e^{\sqrt{\mu_1}t} \left(\sin \alpha + \frac{\cos \alpha}{\sqrt{\mu_1}} \right) + \frac{1}{2} e^{-\sqrt{\mu_1}t} \left(\sin \alpha - \frac{\cos \alpha}{\sqrt{\mu_1}} \right). \end{aligned}$$

Now we use that $\sin \alpha - \cos \alpha / \sqrt{\mu_1} \leq 0$ holds for $\alpha \in [0, \omega_1]$ and conclude

$$0 < u(t; \nu_1(\alpha, \omega_1), \alpha) \leq \frac{1}{2} e^{\sqrt{\mu_1}t} \left(\sin \alpha + \frac{\cos \alpha}{\sqrt{\mu_1}} \right).$$

By explicit calculations it follows that

$$\int_0^{\omega_1} \|u(\cdot; \nu_1(\alpha, \omega_1), \alpha)\|^{-2} d\alpha \geq \frac{4\mu_1}{\exp(2l\sqrt{\mu_1}) - 1}.$$

Similarly, we find

$$\int_0^{\omega_1} \|\tilde{u}(\cdot; \nu_1(0, \beta), \beta)\|^{-2} d\beta \geq \frac{4\mu_1}{\exp(2l\sqrt{\mu_1}) - 1}$$

and (33) implies

$$\mu_1 - \lambda_1 \geq \frac{8\mu_1}{\exp(2l\sqrt{\mu_1}) - 1}. \quad (35)$$

For $l\sqrt{\mu_1} \leq \ln 3$ it follows that $-\lambda_1 \geq 0$. Since the right hand side of (35) is non-increasing the estimate [HLT98]

$$\sqrt{\mu_1} \leq \frac{1}{2} \int_I V(t) dt$$

implies the claimed result. \square

The proof of Theorem 4.1 is an immediate consequence of the results above:

Proof of Theorem 4.1. Using convexity of the map $\lambda \mapsto \lambda^\sigma$ and the Lieb-Thirring inequality (2) we estimate

$$R_\sigma(V; I) = \sum_{k=1}^N \lambda_k^\sigma \leq \sum_{k=1}^N \mu_k^\sigma - (\mu_1^\sigma - \lambda_1^\sigma) \leq L_{\sigma,1}^{\text{cl}} \int_I V(t)^{\sigma+1/2} dt - (\mu_1 - \lambda_1)^\sigma.$$

Hence, for $V \in C_0^\infty(I)$ the claim follows from Lemma 4.4. A standard approximation argument allows us to prove the claim for all non-negative potentials $V \in L^{\sigma+1/2}(I)$. \square

4.2. A sharp Lieb-Thirring inequality with remainder term. Let us now consider general Schrödinger operators H_Ω on bounded or quasi-bounded open sets $\Omega \subset \mathbb{R}^d$ with Dirichlet boundary conditions. To apply the inductive argument introduced in Section 2, fix a coordinate system in \mathbb{R}^d . For $x \in \Omega$ we write $x = (x', t) \in \mathbb{R}^{d-1} \times \mathbb{R}$ and assume that $V_{x'} \in L^{\sigma+d/2}(\Omega(x'))$, a.e. in $x' \in \mathbb{R}^{d-1}$. We use the notation introduced in Section 2 and put

$$A_k(x') = |J_k(x')| \int_{J_k(x')} V_{x'}(t) dt,$$

$$B_k(x') = \int_{J_k(x')} V_{x'}(t) dt.$$

Let $\kappa(x', V) \subset \mathbb{N}$ be the subset of all indices k with $A_k(x') > 2 \ln 3$ and put

$$\Omega_V(x') = \bigcup_{k \in \kappa(x', V)} J_k(x') \subset \mathbb{R} \quad \text{and} \quad \Omega_V = \bigcup_{x' \in \mathbb{R}^{d-1}} \{x'\} \times \Omega_V(x') \subset \Omega.$$

The results from Section 2 and Section 4.1 imply the following sharp Lieb-Thirring inequality with remainder term.

Theorem 4.5. *Let Ω be an open set in \mathbb{R}^d , $d \geq 2$, and assume $\sigma \geq 3/2$. Then the estimate*

$$R_\sigma(V; \Omega) \leq L_{\sigma, d}^{cl} \int_{\Omega_V} V(x)^{\sigma+d/2} dx - L_{\sigma, d-1}^{cl} \int_{\mathbb{R}^{d-1}} \rho(x', V) dx'$$

holds with a remainder

$$\rho(x', V) = \sum_{k \in \kappa(x', V)} \left(\frac{2B_k(x')^2}{\exp(A_k(x')) - 1} \right)^{\sigma+(d-1)/2}.$$

Proof. In view of Proposition 2.1 we have to estimate

$$\text{Tr} W(x', V)^{\sigma+(d-1)/2} = \sum_{k=1}^{N(x')} \text{Tr} H_k(x')_-^{\sigma+(d-1)/2} = \sum_{k=1}^{N(x')} R_{\sigma+(d-1)/2}(V_{x'}; J_k(x')).$$

The potential $V_{x'}$ satisfies the conditions of Theorem 4.1, a.e. in $x' \in \mathbb{R}^{d-1}$. For $k \notin \kappa(x', V)$ we have $|J_k(x')| \int_{J_k(x')} V_{x'} dt \leq 2 \ln 3$ and Theorem 4.1 yields $\text{Tr} H_k(x')_- = 0$. Hence,

$$\begin{aligned} & \text{Tr} W(x', V)^{\sigma+(d-1)/2} \\ &= \sum_{k \in \kappa(x', V)} R_{\sigma+(d-1)/2}(V_{x'}; J_k(x')) \\ &\leq \sum_{k \in \kappa(x', V)} \left(L_{\sigma+(d-1)/2, 1}^{cl} \int_{J_k(x')} V_{x'}(t)^{\sigma+d/2} dt - \left(\frac{2B_k(x')^2}{\exp(A_k(x')) - 1} \right)^{\sigma+(d-1)/2} \right). \end{aligned}$$

Thus the claim follows from Proposition 2.1 using the identities

$$\int_{\mathbb{R}^{d-1}} \sum_{k \in \kappa(x', V)} \int_{J_k(x')} V_{x'}(t)^{\sigma+d/2} dt dx' = \int_{\Omega_V} V(x)^{\sigma+d/2} dx$$

and $L_{\sigma, d-1}^{cl} L_{\sigma+(d-1)/2, 1}^{cl} = L_{\sigma, d}^{cl}$. □

4.3. An example with $V \notin L^{\sigma+d/2}$. Let us illustrate Theorem 4.5 by an example of a Schrödinger operator defined on a horn-shaped region with a potential such that the classical Lieb-Thirring inequality (2) fails. As in Section 3.1 set

$$\Omega_1 = \{(x, y) \in \mathbb{R}^2 : |x| \cdot |y| \leq 1\}$$

and put $V_\alpha(x, y) = |x|^\alpha |y|^{-\alpha}$ with $0 < \alpha < 2/5$. Again, we introduce a scaling parameter $\lambda > 0$ and study the operator

$$H_\alpha = -\Delta - \lambda V_\alpha,$$

defined in $L^2(\Omega_1)$ with Dirichlet boundary conditions. Since $V_\alpha \notin L^{\sigma+1}(\Omega_1)$ the classical results (2) and (1) fail.

Nevertheless, Theorem 4.5 yields an upper bound on $R_\sigma(\lambda V_\alpha; \Omega_1)$ for $3/2 \leq \sigma < (1-\alpha)/\alpha$. Indeed, for any $x \in \mathbb{R}$ the section $\Omega_1(x)$ consists of one open interval $(-x^{-1}, x^{-1})$ and

$$A_1(x) = \frac{4}{|x|} \int_0^{|x|^{-1}} \lambda |x|^\alpha |y|^{-\alpha} dy = \frac{4\lambda}{1-\alpha} |x|^{2(\alpha-1)}.$$

Since $\alpha < 1$ we find that $A_1(x)$ tends to zero as $|x|$ tends to infinity. Thus $A_1(x) \leq 2 \ln 3$ holds for

$$|x| \geq \left(\frac{2\lambda}{(1-\alpha) \ln 3} \right)^{1/(2-2\alpha)} = x_\alpha(\lambda).$$

From Theorem 4.5 it follows that for all $3/2 \leq \sigma < (1-\alpha)/\alpha$ the estimate

$$\begin{aligned} R_\sigma(\lambda V_\alpha; \Omega_1) &\leq 4L_{\sigma,2}^{cl} \int_0^{x_\alpha(\lambda)} \int_0^{x^{-1}} x^{\alpha(\sigma+1)} y^{-\alpha(\sigma+1)} dy dx \lambda^{\sigma+1} \\ &\leq L_{\sigma,2}^{cl} \frac{4}{2\alpha(\sigma+1)(1-\alpha(\sigma+1))} \left(\frac{2}{(1-\alpha) \ln 3} \right)^{\alpha(\sigma+1)/(1-\alpha)} \lambda^{(\sigma+1)/(1-\alpha)} \end{aligned}$$

holds for all $\lambda > 0$.

REFERENCES

- [Ada70] R. A. Adams, *Capacity and compact imbeddings*, J. Math. Mech. **19** (1970), 923–929.
- [AF03] R. A. Adams and J. F. Fournier, *Sobolev spaces*, second ed., Academic Press, 2003.
- [AL78] M. Aizenman and E.H. Lieb, *On semi-classical bounds for eigenvalues of Schrödinger operators*, Phys. Lett. **66** (1978), 427–429.
- [Ber72] F. A. Berezin, *Covariant and contravariant symbols of operators*, Izv. Akad. Nauk SSSR Ser. Mat. **13** (1972), 1134–1167.
- [BS87] M. Sh. Birman and M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space*, Mathematics and its Applications (Soviet Series), D. Reidel Publishing Co., Dordrecht, 1987.
- [CH24] R. Courant and D. Hilbert, *Methoden der mathematischen Physik*, Springer, Berlin, 1924.
- [Cla67] C. Clark, *Rellich's embedding theorem for a 'spiny urchin'*, Canad. Math. Bull **10** (1967), 731–734.
- [DS92] E. B. Davies and B. Simon, *Spectral properties of the Neumann Laplacian of horns*, Geom. Funct. Anal. **2** (1992), 105–117.
- [FG10] R. L. Frank and L. Geisinger, *Two-term spectral asymptotics of the Dirichlet Laplacian on a bounded domain*, to appear in the proceedings of QMath 11, World Scientific, Singapore (2010).
- [Fle78] J. Fleckinger, *Répartition des valeurs propres d'opérateurs elliptiques sur des ouverts non bornés*, C. R. Acad. Sci. Paris Sér. A **286** (1978), no. 3, 149–152.
- [FLU02] J. K. Freericks, E. H. Lieb, and D. Ueltschi, *Segregation in the Falicov-Kimball model*, Comm. Math. Phys. **227** (2002), no. 2, 243–279.
- [GLW11] L. Geisinger, A. Laptev, and T. Weidl, *Geometrical versions of improved Berezin-Li-Yau inequalities*, to appear in the Journal of Spectral Theory (2011).

- [GW10] L. Geisinger and T. Weidl, *Universal bounds for traces of the Dirichlet Laplace operator*, J. Lond. Math. Soc. **82** (2010), no. 2, 395–419.
- [HLT98] D. Hundertmark, E. H. Lieb, and L.E. Thomas, *A sharp bound for an eigenvalue moment of the one-dimensional schrödinger operator*, Adv. Theor. Math. Phys. **2** (1998), 719–731.
- [Hör85] L. Hörmander, *The analysis of linear partial differential operators, vol. 4*, Springer-Verlag, Berlin, 1985.
- [Ivr80] V. Y. Ivrii, *On the second term of the spectral asymptotics for the Laplace-Beltrami operator on manifolds with boundary*, Funktsional. Anal. i Prilozhen. **14** (1980), no. 2, 25–34.
- [Ivr98] ———, *Microlocal analysis and precise spectral asymptotics*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998.
- [KVV09] H. Kovařík, S. Vugalter, and T. Weidl, *Two dimensional Berezin-Li-Yau inequalities with a correction term*, Comm. Math. Phys. **287** (2009), no. 3, 959–981.
- [Lap97] A. Laptev, *Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces*, J. Funct. Anal. **151** (1997), no. 2, 531–545.
- [Lie73] E. H. Lieb, *The classical limit of quantum spin systems*, Comm. Math. Phys. **31** (1973), 327–340.
- [Lie97] ———, *The stability of Matter: From Atoms to Stars, Selecta of Elliot H. Lieb, ed. by W. Thirring*, Springer, 1997.
- [LS10] E. H. Lieb and R. Seiringer, *The stability of matter in quantum mechanics*, Cambridge University Press, Cambridge, 2010.
- [LT76] E. H. Lieb and W. Thirring, *Inequalities for the moments of the eigenvalues of the Schrödinger hamiltonian and their relation to Sobolev inequalities*, Studies in Math. Phys., Essays in Honor of Valentine Bargmann (E. Lieb, B. Simon, and A. S. Wightman, eds.), Princeton Univ. Press, Princeton, New Jersey, 1976, pp. 269–330.
- [Lun10] D. Lundholm, *Weighted supermembrane toy model*, Lett. Math. Phys. **92** (2010), no. 2, 125–141.
- [LW00] A. Laptev and T. Weidl, *Sharp Lieb-Thirring inequalities in high dimensions*, Acta Math. **184** (2000), no. 1, 87–111.
- [LY83] P. Li and S. T. Yau, *On the Schrödinger equation and the eigenvalue problem*, Comm. Math. Phys. **88** (1983), no. 3, 309–318.
- [Mel03] A.D. Melás, *A lower bound for sums of eigenvalues of the Laplacian*, Amer. Math. Soc **131** (2003), 631–636.
- [MM06] S. G. Matinyan and B. Müller, *Adventures of the coupled Yang-Mills oscillators. I. Semiclassical expansion*, J. Phys. A **39** (2006), no. 1, 45–59.
- [RS78] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV. Analysis of Operators*, Academic Press, 1978.
- [Sim83] B. Simon, *Non-classical eigenvalue asymptotics*, J. Functional Anal. **53** (1983), 84–98.
- [SV97] Y. Safarov and D. Vassiliev, *The asymptotic distribution of eigenvalues of partial differential operators*, Translations of Mathematical Monographs, 155, American Mathematical Society, Providence, RI, 1997.
- [vdB84] M. van den Berg, *On the spectrum of the Dirichlet Laplacian for horn-shaped regions in \mathbb{R}^n with infinite volume*, J. Funct. Anal. **58** (1984), 150–156.
- [vdB92a] ———, *Dirichlet-Neumann bracketing for horn-shaped regions*, J. Funct. Anal. **104** (1992), no. 1, 110–120.
- [vdB92b] ———, *On the spectral counting function for the Dirichlet Laplacian*, J. Funct. Anal. **107** (1992), no. 2, 352–361.
- [Wei03] J. Weidmann, *Lineare Operatoren in Hilberträumen. Teil II: Anwendungen*, B.G. Teubner, Stuttgart, 2003.
- [Wei08] T. Weidl, *Improved Berezin-Li-Yau inequalities with a remainder term*, Amer. Math. Soc. Transl. **225** (2008), no. 2, 253–263.
- [Wey12] H. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)*, Math. Ann. **71** (1912), no. 4, 441–479.

Leander Geisinger
Universität Stuttgart
Pfaffenwaldring 57
D - 70569 Stuttgart
E-Mail: geisinger@mathematik.uni-stuttgart.de

Timo Weidl
Universität Stuttgart
Pfaffenwaldring 57
D - 70569 Stuttgart
E-Mail: weidl@mathematik.uni-stuttgart.de

Erschienenene Preprints ab Nummer 2007/001

Komplette Liste: <http://www.mathematik.uni-stuttgart.de/preprints>

- 2011/001 *Geisinger, L.; Weidl, T.*: Sharp spectral estimates in domains of infinite volume
- 2010/018 *Kimmerle, W.; Konovalov, A.*: On integral-like units of modular group rings
- 2010/017 *Gauduchon, P.; Moroianu, A.; Semmelmann, U.*: Almost complex structures on quaternion-Kähler manifolds and inner symmetric spaces
- 2010/016 *Moroianu, A.; Semmelmann, U.*: Clifford structures on Riemannian manifolds
- 2010/015 *Grafarend, E.W.; Kühnel, W.*: A minimal atlas for the rotation group $SO(3)$
- 2010/014 *Weidl, T.*: Semiclassical Spectral Bounds and Beyond
- 2010/013 *Stroppel, M.*: Early explicit examples of non-desarguesian plane geometries
- 2010/012 *Effenberger, F.*: Stacked polytopes and tight triangulations of manifolds
- 2010/011 *Györfi, L.; Walk, H.*: Empirical portfolio selection strategies with proportional transaction costs
- 2010/010 *Kohler, M.; Krzyżak, A.; Walk, H.*: Estimation of the essential supremum of a regression function
- 2010/009 *Geisinger, L.; Laptev, A.; Weidl, T.*: Geometrical Versions of improved Berezin-Li-Yau Inequalities
- 2010/008 *Poppitz, S.; Stroppel, M.*: Polarities of Schellhammer Planes
- 2010/007 *Grundhöfer, T.; Krinn, B.; Stroppel, M.*: Non-existence of isomorphisms between certain unitals
- 2010/006 *Höllig, K.; Hörner, J.; Hoffacker, A.*: Finite Element Analysis with B-Splines: Weighted and Isogeometric Methods
- 2010/005 *Kaltenbacher, B.; Walk, H.*: On convergence of local averaging regression function estimates for the regularization of inverse problems
- 2010/004 *Kühnel, W.; Solanes, G.*: Tight surfaces with boundary
- 2010/003 *Kohler, M.; Walk, H.*: On optimal exercising of American options in discrete time for stationary and ergodic data
- 2010/002 *Gulde, M.; Stroppel, M.*: Stabilizers of Subspaces under Similitudes of the Klein Quadric, and Automorphisms of Heisenberg Algebras
- 2010/001 *Leitner, F.*: Examples of almost Einstein structures on products and in cohomogeneity one
- 2009/008 *Griesemer, M.; Zenk, H.*: On the atomic photoeffect in non-relativistic QED
- 2009/007 *Griesemer, M.; Moeller, J.S.*: Bounds on the minimal energy of translation invariant n-polaron systems
- 2009/006 *Demirel, S.; Harrell II, E.M.*: On semiclassical and universal inequalities for eigenvalues of quantum graphs
- 2009/005 *Bächle, A., Kimmerle, W.*: Torsion subgroups in integral group rings of finite groups
- 2009/004 *Geisinger, L.; Weidl, T.*: Universal bounds for traces of the Dirichlet Laplace operator
- 2009/003 *Walk, H.*: Strong laws of large numbers and nonparametric estimation
- 2009/002 *Leitner, F.*: The collapsing sphere product of Poincaré-Einstein spaces
- 2009/001 *Brehm, U.; Kühnel, W.*: Lattice triangulations of E^3 and of the 3-torus
- 2008/006 *Kohler, M.; Krzyżak, A.; Walk, H.*: Upper bounds for Bermudan options on Markovian data using nonparametric regression and a reduced number of nested Monte Carlo steps
- 2008/005 *Kaltenbacher, B.; Schöpfer, F.; Schuster, T.*: Iterative methods for nonlinear ill-posed problems in Banach spaces: convergence and applications to parameter identification problems
- 2008/004 *Leitner, F.*: Conformally closed Poincaré-Einstein metrics with intersecting scale singularities
- 2008/003 *Effenberger, F.; Kühnel, W.*: Hamiltonian submanifolds of regular polytope

- 2008/002 *Hertweck, M.; Hofert, C.R.; Kimmerle, W.:* Finite groups of units and their composition factors in the integral group rings of the groups $PSL(2, q)$
- 2008/001 *Kovarik, H.; Vugalter, S.; Weidl, T.:* Two dimensional Berezin-Li-Yau inequalities with a correction term
- 2007/006 *Weidl, T.:* Improved Berezin-Li-Yau inequalities with a remainder term
- 2007/005 *Frank, R.L.; Loss, M.; Weidl, T.:* Polya's conjecture in the presence of a constant magnetic field
- 2007/004 *Ekholm, T.; Frank, R.L.; Kovarik, H.:* Eigenvalue estimates for Schrödinger operators on metric trees
- 2007/003 *Lesky, P.H.; Racke, R.:* Elastic and electro-magnetic waves in infinite waveguides
- 2007/002 *Teufel, E.:* Spherical transforms and Radon transforms in Moebius geometry
- 2007/001 *Meister, A.:* Deconvolution from Fourier-oscillating error densities under decay and smoothness restrictions